

LOW-REDSHIFT LY α SELECTED GALAXIES FROM *GALEX* SPECTROSCOPY: A COMPARISON WITH BOTH UV-CONTINUUM SELECTED GALAXIES AND HIGH-REDSHIFT LY α EMITTERS^{1,2,3}

LENNOX L. COWIE,⁴ COWIE@IFA.HAWAII.EDU AMY J. BARGER,^{5,6,4} ESTHER M. HU⁴

Accepted by The Astrophysical Journal

ABSTRACT

We construct a sample of low-redshift Ly α emission-line selected sources from *Galaxy Evolution Explorer* (*GALEX*) grism spectroscopy of nine deep fields to study the role of Ly α emission in galaxy populations with cosmic time. Our final sample consists of 119 (141) sources selected in the redshift interval $z = 0.195 - 0.44$ ($z = 0.65 - 1.25$) from the FUV (NUV) channel. We classify the Ly α sources as active galactic nuclei (AGNs) if high-ionization emission lines are present in their UV spectra and as possible star-forming galaxies otherwise. We classify additional sources as AGNs using line widths for our Ly α emitter (LAE) analysis. These classifications are broadly supported by comparisons with X-ray and optical spectroscopic observations, though the optical spectroscopy identifies a small number of additional AGNs. Defining the *GALEX* LAE sample in the same way as high-redshift LAE samples, we show that LAEs constitute only about 5% of NUV-continuum selected galaxies at $z \sim 0.3$. We also show that they are less common at $z \sim 0.3$ than they are at $z \sim 3$. We find that the $z \sim 0.3$ optically-confirmed Ly α galaxies lie below the metallicity-luminosity relation of the $z \sim 0.3$ NUV-continuum selected galaxies but have similar H α velocity widths at similar luminosities, suggesting that they also lie below the metallicity-mass relation of the NUV-continuum selected galaxies. We show that, on average, the Ly α galaxies have bluer colors, lower extinctions as measured from the Balmer line ratios, and more compact morphologies than the NUV-continuum selected galaxies. Finally, we confirm that the $z \sim 2$ Lyman break galaxies (LBGs) have relatively low metallicities for their luminosities, and we find that they lie in the same metallicity range as the $z \sim 0.3$ Ly α galaxies.

Subject headings: cosmology: observations — galaxies: distances and redshifts — galaxies: abundances — galaxies: evolution — galaxies: starburst

1. INTRODUCTION

High-redshift galaxy studies rely in large part on samples selected using color techniques or extreme emission-line properties. However, it is very difficult to know how such galaxy populations relate to one another or how they fit into the overall scheme of galaxy evolution. Adding complexity to the problem is the fact that active galactic nucleus (AGN) contributions to the light of different galaxy populations may cause different selection biases, often in ways that are hard to determine and quantify. For example, Cowie et al. (2009) showed that at low redshifts the AGN contribution dominates the light at wavelengths below the Lyman continuum edge.

A classic example of a highly valued population whose

relationship to other samples is unclear is the Ly α emitter (LAE) population, which features prominently in the very highest redshift galaxy studies (e.g., Hu & Cowie 2006 and references therein). Current estimates of the fraction of galaxies that exhibit LAE properties are mostly based on comparisons of the LAE and Lyman break galaxy (LBG) populations at $z \sim 2 - 3$ (Shapley et al. 2003). However, it would be useful to know how that fraction varies with redshift, as well as whether the presence of a strong Ly α emission line might be related to other galaxy properties.

Of course, the observed properties of a galaxy are not enough, as one also needs to know how best to translate those properties into interesting physical quantities, such as star formation rates, masses, and metallicities. Because Ly α is resonantly scattered by neutral hydrogen, such translations are particularly difficult to do for LAEs. Determining the escape path of Ly α and hence its dust destruction is an extremely complex problem both theoretically (e.g., Neufeld 1991; Finkelstein et al. 2007) and observationally (e.g., Kunth et al. 2003; Schaerer & Verhamme 2008). In particular, we do not have a very clear understanding of how to translate Ly α luminosities into star formation rates in individual galaxies.

Fortunately, new possibilities for making significant advances in placing the LAEs in context have become available through *Galaxy Evolution Explorer* (*GALEX*; Martin et al. 2005) grism spectroscopy. In a seminal paper Deharveng et al. (2008) have shown that low-redshift ($0.2 < z < 0.35$) galaxies with strong Ly α emission lines can be selected from these data. The selection process

¹ Based in part on data obtained from the Multimission Archive at the Space Telescope Science Institute (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-HST data is provided by the NASA Office of Space Science via grant NAG5-7584 and by other grants and contracts.

² Based in part on data obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the the California Institute of Technology, the University of California, and NASA and was made possible by the generous financial support of the W. M. Keck Foundation.

³ This research used the facilities of the Canadian Astronomy Data Centre operated by the National Research Council of Canada with the support of the Canadian Space Agency.

⁴ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822.

⁵ Department of Astronomy, University of Wisconsin-Madison, 475 North Charter Street, Madison, WI 53706.

⁶ Department of Physics and Astronomy, University of Hawaii, 2505 Correa Road, Honolulu, HI 96822.

is based on finding Ly α in the UV-continuum selected *GALEX* sources and thus is most analogous to locating Ly α emitters in the high-redshift LBG population via spectroscopy (Shapley et al. 2003). However, the procedure enables the selection of a substantial sample of sources that can be compared to the high-redshift LAEs. These low-redshift galaxy samples have many advantages. For one thing, the galaxies are bright and can be easily studied at other wavelengths. Perhaps even more importantly, however, these low-redshift galaxy samples can be integrated into comprehensive studies of the galaxy populations at these redshifts to understand some of the selection biases. For example, we can determine the fraction of NUV-continuum selected galaxies in this redshift interval that have strong Ly α emission. Moreover, by comparing the Ly α properties of the low-redshift galaxies with their optical properties, including their H α line strengths, we can calibrate the conversion of Ly α luminosity to star formation rate and answer many of the questions on how LAEs are drawn from the more general NUV-continuum selected population. Finally, we can see whether the presence of strong Ly α emission in a galaxy is related to the galaxy’s metallicity, extinction, morphology, or kinematics.

The structure of the paper is as follows. In Section 2 we use the *GALEX* extracted spectra (Morrissey et al. 2007) on nine high galactic latitude fields with deep *GALEX* grism exposures to generate a catalog of 261 Ly α emission-line sources in the redshift intervals $z = 0.195 - 0.44$ (low redshift) and $z = 0.65 - 1.25$ (moderate redshift). This sample extends to an NUV (AB) magnitude of 21.8. We divide the Ly α sources into two classes: obvious AGNs, due to the presence of high-excitation lines in their UV spectra, and Galaxies, which do not have such lines. It should be emphasized that some of the objects in the “Galaxy” class may in fact be AGNs, where either the high-excitation lines are weak in the UV or where they lie in missing portions of the UV spectrum.

In Section 3 we present optical spectroscopy of a subsample of the sources with *GALEX* UV spectra. These observations were obtained using multi-object masks with the DEep Imaging Multi-Object Spectrograph (DEIMOS; Faber et al. 2003) on the Keck II 10 m telescope and cover sources both with and without UV spectral identifications.

In Section 4 we use the optical data and a comparison with X-ray observations to show that our AGN and Galaxy classifications from the UV spectra are broadly robust. We also analyze to what extent the *GALEX* line widths may be used to discriminate further between star-forming galaxies and AGNs.

In Section 5 we present the *GALEX* LAE sample defined in the same way as high-redshift LAE samples, and we measure the LAE number counts and luminosity functions. We compare these with the number counts and luminosity functions constructed for both NUV-continuum selected samples and high-redshift LAE samples. We also consider the equivalent width distributions and kinematics of the *GALEX* LAE sample.

In Section 6 we study the properties of low-redshift Ly α galaxies using the Ly α and NUV-continuum selected sources that we optically confirmed as galaxies, as well as an essentially spectroscopically complete sample of

NUV < 24 galaxies in the Great Observatories Origins Deep Survey-North (GOODS-N; Giavalisco et al. 2004) field from Barger et al. (2008). We present the Ly α /H α flux ratios and compare the equivalent widths, metallicities, line widths, colors, extinctions, and morphologies of the galaxies in the two populations.

Finally, in Section 7 we present our conclusions and compare the low-redshift Ly α and NUV-continuum selected galaxy populations with the $z \sim 2$ galaxy population of Erb et al. (2006).

An analysis of the AGN sample will be given in a second paper (A. Barger & L. Cowie 2009, in preparation). Readers who are primarily interested in the results may skip directly to Sections 5, 6, and 7, which are largely self-contained. We use a standard $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M = 0.3$, $\Omega_\Lambda = 0.7$ cosmology throughout.

2. THE GALEX SAMPLE

For the present study we use nine blank high galactic latitude fields with the deepest *GALEX* grism observations. We summarize the fields in Table 1, where we give the *GALEX* name, the J2000 right ascension and declination, the exposure time in kiloseconds, the limiting NUV magnitude to which the spectra were extracted, the galactic $E(B - V)$ in the direction of the field from Schlegel et al. (1998), the galactic latitude, the number of sources with spectra lying within a radius of 32.5 from the field center, and the number of such sources found to have a Ly α emission line in the redshift intervals $z = 0.195 - 0.44$ and $z = 0.65 - 1.25$ based on their UV spectra. The SIRTFFL 00 and SIRTFFL 01 fields have a small overlap. In this region we used the deeper SIRTFFL 00 observations.

For each field we obtained the one and two dimensional spectra of all the sources with *GALEX* grism observations from the Multimission Archive at STScI (MAST). The extracted sources per field constitute nearly complete samples to the NUV limiting magnitudes listed in Table 1. Morrissey et al. (2007) describe the spectral extraction techniques used by the *GALEX* team in analyzing the grism data and detail the properties of the UV spectra.

For our analysis we only consider sources within a 32.5 radius of each field center, since in the outermost regions of the fields there is a higher fraction of poor quality spectra. Even with this restriction the *GALEX* fields are extremely large. In Figure 1 we illustrate the size of a single *GALEX* field (the field around the Hubble Deep Field-North or HDF-N) by showing the positions of the *GALEX* sources with extracted spectra with black squares. We enclose the *GALEX* sources with known redshifts in the literature from Barger et al. (2008) and references therein in a larger open square. We also show the 503 sources found in the deep 2 Ms X-ray observations of the *Chandra* Deep Field-North (CDF-N) from Alexander et al. (2003) with red diamonds and the FUV < 21.5 sources in the GOODS-N with blue triangles.

With our radius restriction we have from just under 300 spectra in the shallowest field to just under 1200 spectra in the deepest field (see Table 1). The FUV spectra cover a wavelength range of approximately 1300 – 1800 Å at a resolution of $\sim 10 \text{ Å}$, and the NUV spectra cover a




FIG. 1.— The positions of the *GALEX* sources with extracted spectra in the field surrounding the HDF-N are shown with small black squares. Those with known redshifts in the literature are enclosed in larger open squares. FUV < 21.5 sources in the GOODS-N are shown with blue triangles. The 503 X-ray sources in the CDF-N are shown with red diamonds. The large black circle shows the 32.5 radius field that we use.

wavelength range of approximately 1850 – 3000 Å at a resolution of ~ 25 Å. However, the spectra become very noisy at the edges of the wavelength ranges. Thus, we only consider sources that have a Ly α emission line in the redshift interval $z = 0.195 - 0.44$ corresponding to 1452.5 – 1750 Å in the FUV spectrum or in the redshift interval $z = 0.65 - 1.25$ corresponding to 2006 – 2735 Å in the NUV spectrum.

We first ran an automatic search procedure for emission lines in the spectra. For each source we fitted the higher signal-to-noise regions in the FUV and NUV spectra separately with a second order polynomial continuum and a Gaussian. Sources with a significant signal in the Gaussian portion were then flagged. We then visually inspected every spectrum to eliminate false emission line detections and to add any cases where the automatic procedure had missed an emission line. Most of the emission lines were detected by the automatic procedure with only a relatively small number of corrections required from the visual check. A very small number of sources were eliminated based on visual inspection of the images. On average 4% of the sources (recall that these are NUV-continuum selected and comprise both galaxies and a significant number of stars) have detected emission lines. This percentage is relatively invariant from field to field (see Table 1), though it is slightly higher in the deeper fields; in the four deepest fields, 5% of the sources have detected emission lines.

We next measured a redshift for each source with an emission line and split the sources into one of two classes: an AGN class if there were high-excitation lines (usually OVI or CIV) present in addition to the Ly α line, or a Galaxy class to denote a potential star-forming galaxy if




FIG. 2.— Some sample *GALEX* spectra illustrating the classifications for the Ly α selected sources. (a) Broad-line AGN. The positions of the various broad emission lines are marked, as is the position of a strong intervening Lyman limit system. (b) Narrower line AGN, where strong CIV and NV (not marked) are seen in addition to the Ly α line. (c) A spectrum classified as a candidate Ly α Galaxy. There is no emission seen at the CIV position, and the single line is narrow.

there were only a single line visible. In the latter case we assumed the line was Ly α in determining the redshift, an assumption we will test and find to be extremely reliable using optical spectra. We will frequently refer to the sources in our Galaxy class as our candidate Ly α Galaxies, using the word “candidate” to emphasize that




FIG. 3.— The rest-frame $\text{EW}(\text{Ly}\alpha)$ distribution for our candidate $\text{Ly}\alpha$ Galaxy sample. The numbers correspond to 4 Å bins. The red line shows the rest-frame $\text{EW}(\text{Ly}\alpha) \sim 15$ Å limit of the sample. The blue curve shows the exponential fit to the data described in the text, which has a scale length of 23 Å.

some of these will turn out to be AGNs. The relative classifications are illustrated in Figure 2, where we show examples of (a) a broad-line AGN, (b) a narrower line AGN, and (c) a candidate $\text{Ly}\alpha$ Galaxy.

For each source we fitted a 140 Å rest-frame region around the $\text{Ly}\alpha$ line with a Gaussian and a flat continuum to determine four parameters: the $\text{Ly}\alpha$ line width (this includes the rather wide instrumental resolution), the observed-frame equivalent width (EW) of the line, the central wavelength, and the continuum level. We used the IDL MPFIT procedures of Markwardt (2008), which are based on the More (1978) implementation of the Levenberg-Marquardt algorithm and are very robust. We used the *GALEX* noise vector to determine the input error for the spectrum and determined the statistical errors from the covariance matrix returned by the program.

All of the sources have detectable continuum at the $\text{Ly}\alpha$ position. This is a direct consequence of the NUV magnitude selection in the *GALEX* spectral extraction for the NUV selected $\text{Ly}\alpha$ lines, and of the NUV magnitude selection and the small difference between the FUV and NUV magnitudes for the FUV selected $\text{Ly}\alpha$ lines at lower redshifts. However, changes in the fitting procedure (such as choosing a different wavelength range or fitting the baseline with a linear fit rather than a constant) can change the fitted parameters. In general these changes correspond to errors in the EW of less than 20%, and, for the most part, the errors are smaller than 10%, which are comparable to the statistical errors. However, in a very small number of cases (less than a few percent) more substantial errors can occur as the fitting procedure finds a substantially different solution. Visual inspection suggests that the present choice of parameters provides the most reasonable results.

The final $\text{Ly}\alpha$ selected samples are summarized in Tables 2–10 for each of the nine fields individually. For each source we give the J2000 right ascension and declination, the NUV and FUV magnitudes, the redshift, the

line width in km s^{-1} together with the 1σ error, and the AGN or Galaxy classification from the UV spectra. The NUV and FUV magnitudes are from the deep broadband *GALEX* images of the fields with exposure times ranging from 26 ks to 240 ks in the NUV band and 28 ks to 120 ks in the FUV band. For the shallowest exposures the 1σ error for the NUV band is approximately 26, and for the FUV band it is approximately 26.4. The sources with a Galaxy classification can be compared with those given in Deharveng et al. (2008) for the subsample of the present fields analyzed in that work. In general the agreement is very good. The small number of sources omitted from one list but included in the other are generally marginal cases where the inclusion or exclusion is somewhat arbitrary. We provide a more detailed comparison in Section 5.1.

We show the distribution of the rest-frame $\text{EW}(\text{Ly}\alpha)$ for our candidate $\text{Ly}\alpha$ Galaxies in Figure 3. For the $z = 0.195 - 0.44$ redshift interval the sample appears to be relatively complete down to a rest-frame $\text{EW}(\text{Ly}\alpha) \sim 15$ Å, which is shown by the red line. (More precisely, the limit of the observed-frame $\text{EW}(\text{Ly}\alpha)$ is ~ 20 Å.) This limit is about a factor of two times higher in the $z = 0.65 - 1.25$ redshift interval. In our comparisons with high-redshift LAE samples in Section 5, we will include only sources having a rest-frame $\text{EW}(\text{Ly}\alpha) > 20$ Å, which is normally used as the definition of high-redshift LAEs (e.g., Hu et al. 1998). Such a sample should be very robust for the $z = 0.195 - 0.44$ interval. The blue curve shows an exponential fit of the form $\exp(-\text{EW}/\text{scale length})$ to the data above a rest-frame EW of 20 Å where we believe the sample to be substantially complete. The normalization is 29 ± 4 , and the EW scale length is 23.7 ± 2.2 Å. This distribution is nearly identical in shape to that found by Shapley et al. (2003) for the $z \sim 3$ LBGs.

Only seven of our candidate $\text{Ly}\alpha$ Galaxies have a rest-frame $\text{EW}(\text{Ly}\alpha) > 100$ Å. As can be seen from Figure 3, five of these lie in the 100 – 120 Å range. There are two further sources which lie above the upper bound of the figure, but these are all almost certainly broad-line AGN where we are only seeing the $\text{Ly}\alpha$ line.

In Figure 4 we show the redshift distributions of our AGNs (red squares) and our candidate $\text{Ly}\alpha$ Galaxies (black diamonds) versus magnitude. In (a) we plot the sources versus NUV magnitude and in (b) versus FUV magnitude. Nearly all of the more luminous sources are AGNs. As Deharveng et al. (2008) note, this results in there being relatively few candidate $\text{Ly}\alpha$ Galaxies in the moderate-redshift interval. In the low-redshift interval the candidate $\text{Ly}\alpha$ Galaxy population begins to enter in substantial numbers around a NUV magnitude of 21.

Figure 4(b) suggests immediately that we are misclassifying some sources as candidate $\text{Ly}\alpha$ Galaxies when they are in fact AGNs, since we would expect $z \sim 1$ galaxies to be very faint or undetected in the FUV band, which lies below the Lyman continuum edge at these redshifts (e.g., Siana et al. 2007; Cowie et al. 2009). Quantifying this contamination is one of the keys to understanding the star-forming population. We will address this issue in Section 4.

We will require the observed area as a function of NUV magnitude when we calculate the number counts and luminosity functions of the LAEs. We determined this for




FIG. 4.— (a) Redshift vs. NUV magnitude for our Ly α selected sample from all nine *GALEX* fields. Sources classified as AGNs based on high-excitation lines in the spectra (otherwise) are shown as red squares (black diamonds). The depths of the fields range from NUV(AB) = 20.5 in HDFN 00 to 21.8 in GROTH 00. (b) Redshift vs. FUV magnitude with symbols as in (a). Here sources fainter than 24.7 are shown at a nominal magnitude of 24.7. (c) Total observed area for the nine fields vs. NUV magnitude.

each field separately by computing the ratio of sources with *GALEX* spectra (identified or not) at a given NUV magnitude to sources with that NUV magnitude in the continuum catalog. We then multiplied this ratio for each field by the area corresponding to the 32'5 selection radius and summed the results for the nine fields to

form the area-magnitude relation shown in Figure 4(c). At magnitudes brighter than NUV = 20.5 the area is the 8.2 deg² area of the nine fields. The area then drops as we reach the limiting magnitudes of each of the individual fields, eventually falling to zero at ~ 21.8 , which is the limiting depth of the deepest (GROTH 00) field.

Given the low spatial resolution of the *GALEX* data, some sources may be blends. While this does not affect the Ly α fluxes or luminosities, it will diminish the EWs if the measured continuum is too high relative to the line due to a blend. It will also boost the NUV magnitudes so that we are including sources which are fainter than the nominal limits.

We first re-measured the *GALEX* coordinates by visually determining the peak position in the NUV image closest to the cataloged position in the data archive. It is these coordinates which we use in the optical spectroscopy. The dispersion of the re-measured positions relative to the catalog positions is 0''.5. However, a small number of sources have more significant offsets: about 6% have offsets greater than 1''.5 and 2% greater than 3''. In Tables 2 through 10 we show our re-measured coordinates, marking those which have a significant offset from the cataloged position in the data archive with the label (o) in the name column.

We next tested for the fraction of blends by comparing the NUV images with the ultradeep u^* -band images obtained on the GROTH 00 and COSMOS 00 fields as part of the Canada-France-Hawaii Telescope (CFHT) Legacy Survey deep observations. These are available from the MegaPipe database at the Canadian Astronomy Data Centre (CADC). We corrected for the small absolute astrometric offsets between the CFHT and *GALEX* coordinates prior to the comparison. We find that about 5% are significant enough blends to seriously affect the photometry and coordinates. We mark these in Tables 2 and 5 with a (b) in the name column. The blending problem is small enough to be neglected in the analysis, but the small number of objects with significant positional offsets should be borne in mind when using the *GALEX* coordinates in the data archive.

3. OPTICAL SPECTROSCOPY

While a number of sources in our *GALEX* AGN and candidate Ly α Galaxy samples have pre-existing optical redshifts, these identifications are generally selected in some way, often from optical colors or because of the presence of X-ray emission or other AGN signatures. However, in order to understand the statistical distribution of the optical properties of the sources in these samples, it is critical to observe at optical wavelengths sources randomly chosen from these samples.

In addition, in order to understand how the *GALEX* Ly α selected sources relate to the more general population of NUV-continuum selected sources, it is important to observe at optical wavelengths sources randomly chosen from the *GALEX* spectroscopic sample without UV spectral identifications (i.e., without strong Ly α emission).

We therefore made optical spectroscopic follow-up observations with the DEIMOS spectrograph (Faber et al. 2003) on the Keck II 10 m telescope. We obtained observations of 39 sources randomly chosen from our *GALEX* candidate Ly α Galaxy sample, 38 of which lie in the red-

shift interval $z = 0.195 - 0.44$ and one in the redshift interval $z = 0.65 - 1.25$ (Table 11), 15 sources randomly chosen from our *GALEX* AGN sample (Table 12), and 124 sources randomly chosen from our *GALEX* spectroscopic sample without UV spectral identifications. For the latter we summarize only the 31 sources found to lie in the redshift interval $z = 0.195 - 0.44$ (Table 13).

In Tables 11–13 we give the *GALEX* source name, the J2000 right ascension and declination, the NUV magnitude, the SDSS model C g magnitude from the DR6 release (Adelman-McCarthy et al. 2008; these are only given in Tables 11 and 13), the *GALEX* UV redshift (these are only given in Tables 11 and 12), the optical spectroscopic redshift, the observed-frame $\text{EW}(\text{H}\alpha)$ in $\text{\AA}$ and its 1σ error (these are only given in Tables 11 and 13), and the spectral type that we determined from the line widths of the optical spectrum.

In order to provide the widest possible wavelength coverage, we used the ZD600 ℓ/mm grating. We used $1''$ wide slitlets, which in this configuration give a resolution of $4.5 \text{ \AA}$, sufficient to distinguish the $[\text{O II}]\lambda 3727$ doublet structure for the higher redshift sources ($z > 0.65$). The spectra cover a wavelength range of approximately $5000 \text{ \AA}$ and were centered at an average wavelength of $6800 \text{ \AA}$, though the exact wavelength range for each spectrum depends on the slit position with respect to the center of the mask along the dispersion direction. Each $\sim 0.5 \text{ hr}$ exposure was broken into three subsets with the objects stepped along the slit by $1''.5 - 2''$ in each direction. The two-dimensional spectra were reduced following the procedure described in Cowie et al. (1996), and the final one-dimensional spectra were extracted using a profile weighting based on the continuum of the spectrum. Calibration stars were observed in the same configuration on each of the individual nights.

The spectra were all obtained at a near-parallactic angle to minimize atmospheric refraction effects. Each spectrum was initially calibrated using the measured response from a calibration star. However, since some of the spectra were obtained during a night of varying extinction, the absolute flux calibration using calibration stars is sometimes problematic. We also note that relative slit losses always pose a problem, even for the photometric nights. We show pieces of the spectra of two of the *GALEX* Ly α candidate Galaxies in Figures 5 and 6. The absolute fluxes are based on the calibration star, but the optical spectrum in Figure 5 was not observed in photometric conditions.

An extensive discussion of the various procedures that may be used to determine line fluxes from this type of spectrum can be found in Kakazu et al. (2007). However, here we focus only on the measurements of the relative line fluxes, which we need for the metallicity diagnostics, and on the measurements of the absolute line fluxes, which we require for the determination of the Ly α /H α flux ratio.

For each spectrum we fitted a standard set of lines. For the stronger lines we used a full Gaussian fit together with a linear fit to the continuum baseline. For weaker lines we held the full width constant using the value measured in the stronger lines and set the central wavelength to the nominal redshifted value. We fitted the $[\text{O II}]\lambda 3727$ line with two Gaussians with the appropriate wavelength




FIG. 5.— The optical and UV spectrum for the candidate Ly α Galaxy GALEX1417 + 5228 in the GROTH 00 field at $z = 0.206$ (see Table 11). The optical spectrum is dominated by the H α line (in a) and the $[\text{O III}]\lambda 3727$ doublet (in b). (c) The H γ and $[\text{O III}]\lambda 4363$ auroral lines in the spectrum. (d) The UV spectrum. There are no signs of any broad lines in the optical spectrum, and the line ratios are consistent with this being an extremely low-metallicity emission-line galaxy. The optical spectrum panels are shown in μJy units, but since the absolute calibration is not well determined due to not having photometric conditions, we have not labeled them.




FIG. 6.— The (a) optical and (b) UV spectrum for the candidate Ly α Galaxy GALEX1047+5827 in the LOCK 00 field at $z = 0.2436$ (see Table 11). From the optical data we classify this source as a type 1.8 Seyfert galaxy based on the broad underlying H α line in the optical spectrum and a corresponding weak broad H β line. The optical spectrum was taken in photometric conditions. While no high-excitation lines are seen in the UV spectrum, the Ly α line in the UV spectrum is broader than the instrumental resolution.

separation. We also measured the noise as a function of wavelength by fitting Gaussians with the measured line-width from the strong lines at random positions in the spectrum and computing the dispersion in the results. Once again systematic errors from the choice of fitting procedure and wavelength range may exceed the statistical errors and can be as large as 10 – 20% of the EW.

As long as we restrict the line measurements to short wavelength ranges where the DEIMOS response is essentially constant, we can robustly measure the ratios of line fluxes from the spectra without any flux calibration. For example, we can assume that the responses of neighboring lines (e.g., H β and [O III] λ 5007) are the same and therefore measure their flux ratios without calibration. In this regard the problems of distinguishing AGNs from emission-line galaxies using Baldwin et al. (1981) diagnostic diagrams and of determining metallicities is considerably simplified by the presence of H α near [NII] λ 6584 (Figure 5(a)), H β near [O III] λ 4949 and [O III] λ 5007 (Figure 5(b)), and H γ near [O III] λ 4363 (Figure 5(c)). For example, by assuming case B ratios,

we can take the ratios of the Balmer lines accompanying the metal lines and the ratios of the metal lines to determine the extinction-corrected metal line ratios. Unfortunately, we cannot so easily do this near [O II] λ 3727, where the Balmer lines are weak and in some cases contaminated. Here we must rely on the flux calibration made using the measured response from the calibration star. For this reason we will use the [NII] λ 6584/H α ratio as our primary metal diagnostic.

In order to measure the H α line flux, we determined the continuum flux from the SDSS model C magnitude (DR6 release; Adelman-McCarthy et al. 2008) at the SDSS bandpass closest to redshifted H α , and we multiplied this by the observed-frame EW(H α). This is an approximation, since it assumes that the measured EW(H α) is representative of the value averaged over the total light of the galaxy, including regions outside the slit. However, for the photometric cases we derive crudely similar values directly from the calibrated spectra, suggesting the procedure is relatively robust. In one case (GALEX1417+5228) there is an H α flux measurement of $1.3 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ from Liang et al. (2004), which may be compared with the presently derived value of $1.9 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$. The Liang et al. value is based on calibrated slit spectra and may be an underestimate, since they measure a corresponding H β flux of $0.61 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$, which for the case B ratio would give an H α flux of $1.7 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ in the absence of any extinction correction and a higher value if any extinction were present. However, we will assume, based on this comparison, that systematic errors in the absolute H α flux levels could be as high as 50%.

4. AGN-GALAXY DISCRIMINATION

We may expect that sources classified as AGNs based on the presence of high-excitation lines in the UV spectra are truly AGNs. However, the classification of the candidate Ly α Galaxies is not so clean, and they may contain a substantial amount of AGN contamination. In some cases the high-excitation lines of these AGNs may fall at problem wavelengths, and in other cases the high-excitation lines may be intrinsically weak. The optical spectroscopic data support these points. All but one of the 15 AGNs classified as such from the UV spectra and then observed in the optical are straightforwardly identified as intermediate-type Seyfert galaxies (see Table 12; since only one source appears as an emission-line galaxy, this suggests that for the most part one would know that these sources are AGNs based solely on their optical spectra). However, three of the 39 candidate Ly α Galaxies classified as such from the UV spectra and then observed in the optical are also identified as intermediate-type Seyfert galaxies (see Table 11 and, e.g., Figure 6). The remaining 36 do not have visible broad lines (see, e.g., Figure 5). This suggests that we have about an 8% broad-line AGN contamination rate in this population. Finally, we note that the 31 sources chosen randomly from the GALEX spectroscopic sample without UV spectral identifications that happen to lie in the redshift interval $z = 0.195 - 0.44$ all appear to be star-forming galaxies with H α luminosities comparable to the candidate Ly α Galaxies (see Table 13).

There are deep X-ray observations of portions of the GALEX areas for several of the fields used in our study.




FIG. 7.— Comparison of the *GALEX* Ly α selected sources with the *Chandra* X-ray sources in the (a) HDFN 00 (CDF-N), (b) LOCK 00 (CLANS), and (c) CDFS 00 (CDF-S and Extended CDF-S) fields. The X-ray sources are shown with red solid diamonds, except in the CDFS 00 field where the 2 Ms CDF-S (Extended CDF-S) X-ray sources are shown with purple (red) solid diamonds. The *GALEX* sources are shown with black (blue) open squares for AGNs based on the presence of high-excitation lines in their UV spectra (otherwise).

In Figure 7 we illustrate the overlap of the X-ray sources (red solid diamonds) with the *GALEX* Ly α selected sources (black open squares for AGNs based on their UV spectra; blue open squares otherwise). In (a) we compare the CDF-N sources (Alexander et al. 2003) with the HDFN 00 sources; in (b) we compare the CLANS sources (Trouille et al. 2008, 2009) with the LOCK 00 sources; and in (c) we compare the Extended CDF-S sources (Lehmer et al. 2005; Virani et al. 2006) and the 2 Ms CDF-S sources (purple solid diamonds; Luo et al. 2008) with the CDFS 00 sources. In total eleven sources in the *GALEX* Ly α selected sample lie in the X-ray fields. We summarize their X-ray properties in Table 14, where we give the *GALEX* name of the source, the J2000 right ascension and declination, the 2 – 8 keV and 0.5 – 2 keV fluxes, the 2 – 8 keV luminosity computed using the procedure given in Barger et al. (2005), the redshift from the UV spectra, and the classification of the source as an AGN or a candidate Ly α Galaxy from the UV spectrum.

All nine of the sources classified as AGNs based on their UV spectra are also X-ray sources and have logarithmic 2 – 8 keV luminosities in the range $\log L_X = 43.24 - 44.55 \text{ erg s}^{-1}$ (see Table 14), placing them at or near quasar luminosities ($\log L_X > 44$) and near the break in the AGN luminosity function at $z \lesssim 1.2$ (e.g., Yencho et al. 2009). In contrast, of the two candidate Ly α Galaxies, one is only weakly detected in the 0.5 – 2 keV band in the extremely deep 2 Ms CDF-S image, and the other is not detected in the Extended CDF-S image. This places both of their $\log L_X$ values below 41 erg s^{-1} (see Table 14), suggesting that they are strong star-forming galaxies rather than AGNs. Thus, in all of the overlapped sources, the UV classification is robustly supported by the X-ray data.

We can also make a comparison with a previous optical spectroscopic sample, since Prescott et al. (2006) observed a large sample of candidate AGNs in a 1.39 deg^2 region of the COSMOS *HST* Treasury field (Scoville et al. 2007), which overlaps significantly with the *GALEX* COSMOS 00 field. In Figure 8 we plot the Ly α line widths for the *GALEX* Ly α selected sample (red solid squares for AGNs based on their UV spectra; black small diamonds otherwise) that lie in the overlap region with the Prescott et al. (2006) observations versus redshift. Where Prescott et al. (2006) classifications exist for these sources, we show them on the figure (black large open squares for optically selected AGNs; blue solid diamonds for optical emission-line galaxies). All eleven of the *GALEX* AGNs with a Prescott et al. (2006) optical identification were classified by those authors as AGNs based on their optical spectra, and all three of the *GALEX* candidate Ly α Galaxies with a Prescott et al. (2006) optical identification were classified by those authors as emission-line galaxies based on their optical spectra. All of the redshifts in the two samples are fully consistent.

As Figure 8 illustrates, the Ly α line widths provide a second diagnostic that may be used to separate AGNs from emission-line Galaxies. In general, at the low resolution of the *GALEX* grisms the emission lines from galaxies should be essentially unresolved, and only broad AGN lines ($\sim$ several thousand km s^{-1}) should produce significant broadening in the *GALEX* spectra. (In the present




FIG. 8.— Comparison of the rest-frame line widths of the *GALEX* Ly α selected sample in the COSMOS 00 field with those of optically selected AGNs in the COSMOS field from Prescott et al. (2006) vs. redshift. The *GALEX* sources are plotted with red solid squares (AGNs based on the presence of high-excitation lines in their UV spectra) and black small diamonds (otherwise). The error bars show the $\pm 1\sigma$ range in the line widths. The *GALEX* sources with matches to the optically selected AGNs in Prescott et al. are surrounded by black large open squares. The *GALEX* sources with matches to the emission-line galaxies in Prescott et al. are shown with blue solid diamonds. The blue curves show the varying instrumental resolution for a point source.




FIG. 9.— Rest-frame line widths of the *GALEX* Ly α selected sample in the low-redshift interval. The sources are plotted with red solid squares (AGNs based on the presence of high-excitation lines in their UV spectra) and black diamonds (otherwise). The error bars show the $\pm 1\sigma$ range in the line widths. The blue curve shows the varying instrumental resolution for a point source.

paper we only determine the intrinsic galaxy line widths from the optical spectroscopy.) This problem is further complicated by the nature of the grism data where the final resolution depends on the image size. Deharveng et al. (2008) used the line width as a second criterion to separate AGNs from emission-line Galaxies in their *GALEX* sample. However, as we illustrate further in Figure 9, a good deal of scatter in the measured line widths and a substantial overlap between AGNs and candidate Ly α Galaxies make the choice of a dividing line difficult. In particular, the average line width of the candidate Ly α Galaxies appears to be larger than the nominal resolution




FIG. 10.— Distribution of observed-frame line widths in the (a) $z = 0.195 - 0.44$ and (b) $z = 0.65 - 1.25$ *GALEX* Ly α selected samples. The black open histogram shows the candidate Ly α Galaxies, and the red open histogram shows the AGNs (vertically displaced by 20 in (a) and 10 in (b) for clarity). In each panel the blue vertical lines show the range of the instrumental resolution in the wavelength interval, and the black vertical dashed line shows the proposed split above which we consider that the FWHM line width alone may show a source to be an AGN (i.e., 15 Å in the FUV spectra and 30 Å in the NUV spectra). In (a) the blue (red) shaded histogram shows sources in the candidate Ly α Galaxy population confirmed to be Galaxies (broad-line AGNs) from optical spectroscopy. In (b) the cyan shaded histogram shows sources in the candidate Ly α Galaxy population with strong Lyman continuum breaks.

of a point source in the *GALEX* data.

We consider this more quantitatively in Figure 10(a), where we show the distribution of line widths (now in the observed frame so we may compare with the instrumental resolution) in the *GALEX* $z = 0.195 - 0.44$ Ly α selected sample. We show the distribution for the candidate Ly α Galaxies with the black open histogram, and we show the distribution for the sources classified as AGNs based on their UV spectra with the red open histogram (offset upwards for clarity). There is a substantial spread of the Ly α Galaxies about the instrumental resolution (whose range with wavelength is shown by the blue vertical lines) resulting from the effects of statistical and systematic noise in the Gaussian fitting. As we have discussed above, the lines are also wider on average than the nominal instrumental resolution for a point-like object.




FIG. 11.— Comparison of the broadband FUV–NUV colors of the *GALEX* candidate Ly α Galaxies that satisfy the line width cuts (black squares) with the colors of a spectroscopically complete sample of NUV < 23 galaxies in the GOODS-N from Barger et al. (2008; red diamonds when identified as AGNs based on their X-ray luminosities and blue diamonds otherwise). Note that one of the *GALEX* $z = 0.65 - 1.25$ candidate Ly α Galaxies has a color redder than the y -axis upper limit of the plot. Sources with FUV > 25.2 are shown at that value with upward-pointing arrows. For most of the data points the errors in the colors are small compared to the symbol sizes.

We show all 33 candidate Ly α Galaxies that we have confirmed as star-forming emission-line galaxies using our DEIMOS optical spectroscopy (see Table 11; hereafter, we refer to these as our optically-confirmed Ly α Galaxies) with the blue shaded histogram. This excludes the four galaxies with broad lines in the optical spectra labeled as AGNs in Table 11, as well as one additional object classified as an AGN based on its optical line ratios (see Section 6.3). All of the optically-confirmed Ly α Galaxies lie near the instrumental resolution, as do two of the four candidate Ly α Galaxies that we found to be broad-line AGNs using our DEIMOS optical spectroscopy (see Table 11; red shaded histogram). The remaining two AGNs (including the one in the LOCK 00 field shown in Figure 6) have substantial Ly α line widths. We have chosen an observed-frame FWHM line width of 15 Å, shown as the black vertical dashed line in Figure 10(a), to roughly split the spread of true galaxy line widths from the extended tail, which may be predominantly AGN-dominated. However, further work is clearly needed to better define this separation, and, even with this elimination, there will be AGNs where the lines are not resolved (as is evident from the red open histogram). There are 72 candidate Ly α Galaxies in Figure 10(a) with widths less than 15 Å, of which 33 are optically-confirmed Ly α Galaxies and 3 are optically-confirmed AGNs (this includes the AGN identified from the line ratios). This suggests that even with a line width criterion, we have about a 10% AGN contamination in the candidate Ly α Galaxy sample.

In Figure 10(b) we show the distribution of Ly α line widths in the *GALEX* $z = 0.65 - 1.25$ Ly α selected sample. Here most of the candidate Ly α Galaxies are clearly too broad relative to the instrumental resolution (whose range with wavelength is shown by the blue verti-

cal lines), even when allowance is made for the line-width noise and the uncertainty in the instrumental resolution, and must be AGNs. We consider only the 6 sources with observed-frame FWHM line widths less than 30 Å (black vertical dashed line) to be plausible candidate Ly α Galaxies. However, in contrast to the lower redshift sources, we can test this classification further by looking for the presence of the Lyman break at the Lyman continuum edge. The nine sources with possible Lyman continuum breaks are shown by the cyan shaded histogram in Figure 10(b). Four of the six sources in the low-velocity width range (i.e., < 30 Å) have breaks, suggesting that at least for these sources we have confirmation of the line width classification by the Lyman break criterion. Alternatively, however, they may just be type 2 AGNs with Lyman continuum breaks, as the > 30 Å sources with Lyman continuum breaks likely are. Only one of these sources (Table 11) has been observed and confirmed as a star former with optical spectroscopy.

We test our line width cuts in Figure 11, where we compare the colors of the candidate Ly α Galaxies that satisfy our line width cuts (black squares) with the colors of a spectroscopically complete sample of NUV < 23 galaxies in the GOODS-N from Barger et al. (2008; red diamonds for sources classified as AGNs based on having X-ray luminosities > 10⁴² erg s⁻¹ and blue diamonds otherwise). Sources fainter than FUV = 25.2 are shown at this value with upward-pointing arrows. The *GALEX* candidate Ly α Galaxies generally lie along the color track of the GOODS-N galaxies, except for the two $z = 0.65 - 1.25$ sources without Lyman breaks whose colors are well below the track, placing them in the AGN category. As can also be seen from Figure 11, many of the AGNs in the GOODS-N follow the galaxy color track, so the *GALEX* Ly α Galaxy sample might have further AGN contamination. Thus, the four sources with a Lyman continuum break and an observed-frame line width less than 30 Å in the $z = 0.65 - 1.25$ interval should be considered as an upper limit on the Ly α selected star-forming galaxy population in this redshift interval.

5. COMPARISON OF THE GALEX LAE SAMPLE WITH HIGH-REDSHIFT LAE SAMPLES

5.1. LAE Selection

In order to make valid comparisons with the high-redshift LAE samples, we need to construct our *GALEX* LAE sample carefully. We start with our candidate Ly α Galaxy sample, which already excludes sources with detectable high-excitation lines in the UV spectra. We now also exclude sources with observed-frame FWHM line widths greater than 15 Å (30 Å) in the redshift interval $z = 0.195 - 0.44$ ($z = 0.65 - 1.25$). For the moderate-redshift interval we further exclude the two < 30 Å sources without a strong Lyman continuum break.

In Figure 12 we plot rest-frame EW(Ly α) versus redshift for the above sample. A substantial fraction of the sources have an EW(Ly α) less than or equal to the 20 Å value normally used to define the high-redshift LAE population (e.g., Hu et al. 1998). If we follow the EW(Ly α) > 20 Å definition, we are left with 41 LAEs in the low-redshift interval and 4 in the moderate-redshift interval. This LAE sample is tabulated in Table 15 (low-




FIG. 12.— Rest-frame $\text{EW}(\text{Ly}\alpha)$ for the candidate $\text{Ly}\alpha$ Galaxies vs. redshift. This consists of objects with no high-excitation lines in the UV and with widths less than 15 Å in the low-redshift interval and less than 30 Å in the moderate-redshift interval. Objects with $\text{FUV-NUV} < 1.8$ are also excluded in the moderate-redshift interval. The final LAE sample is taken to be those sources with a rest-frame $\text{EW}(\text{Ly}\alpha) > 20$ Å (red dashed line). The error bars show the $\pm 1\sigma$ errors from the formal statistical fit of the Gaussian and baseline. As discussed in the text, there may be comparable systematic errors resulting from the choice of fitting procedure.

redshift interval) and Table 16 (moderate-redshift interval), where we give the name, the J2000 right ascension and declination, the NUV and FUV magnitudes, the *GALEX* redshift, the logarithm of the $\text{Ly}\alpha$ luminosity, $L = 4\pi d_L^2 f(\text{Ly}\alpha)$, and its 1σ error, where $f(\text{Ly}\alpha)$ is the observed line flux and d_L is the luminosity distance, the rest-frame $\text{EW}(\text{Ly}\alpha)$ and its 1σ error, and the ground-based redshift, if available. The tables are sorted by rest-frame $\text{EW}(\text{Ly}\alpha)$, with the highest EW first. We also include in the table sources with $15 \text{ Å} < \text{EW}(\text{Ly}\alpha) \leq 20 \text{ Å}$, though these are not part of our final LAE sample.

The ground-based redshifts are based on this paper’s observations, except for two sources in the COSMOS 00 field, where the redshifts are from Prescott et al. (2006), and two sources in the GROTH 00 field, where the redshifts are from Finkelstein et al. (2009). The redshifts for the Prescott et al. and Finkelstein et al. sources are enclosed in parentheses and annotated with a “pr” or “f”. Hereafter, we concentrate on the low-redshift LAE sample, since the moderate-redshift LAE sample is so small.

The low-redshift sample may be expected to be relatively complete, but, as we have discussed, we expect it to have $\sim 20\%$ AGN contamination. For our LAE analysis we do not eliminate any sources that we know to be AGNs from their optical spectra, since we do not have this information for a substantial part of the sample. However, we do note in the table where a source is clearly an AGN based on its optical spectrum, either because it has broad lines or based on the Baldwin et al. (1981) diagnostic diagram of $[\text{OIII}]\lambda 5007/\text{H}\beta$ versus $[\text{NII}]\lambda 6584/\text{H}\alpha$. We denote these latter sources with BPT AGN. Finkelstein et al. (2009) classify the two sources in the GROTH 00 field where we have used their redshifts as AGNs. In one case this is based on broad lines, and in the other it is based on the

Baldwin et al. (1981) diagnostic. They also classified GALEX1417+5228 as an AGN, but we believe this source to be a very high-excitation star former. We discuss this very interesting source in Section 6.3.

Considering only the sources for which we have obtained optical spectroscopic data and classifying sources as AGNs only if they show AGN signatures in their optical spectra, we find the AGN contamination in our final LAE sample to be 4 out of 23 or $9 - 31\%$, where the range is $\pm 1\sigma$. Finkelstein et al. (2009) give $17 - 61\%$ for this range based on observations of a subsample of the Deharvang et al. (2008) sources in the GROTH 00 field. The small sample sizes leave the exact value somewhat statistically uncertain, but it is important to note that the degree of remaining AGN contamination is a function of the NUV-continuum selection procedure and may vary with different samples, so it is not a particularly interesting or physical quantity in itself. The more interesting quantity is the total fraction of AGNs in all the NUV-continuum selected sources in a given redshift and magnitude range, including both those identified in the UV and the additional objects identified with the optical spectra. We shall return to analyze this further in A. Barger & L. Cowie (2010, in preparation).

Our low-redshift LAE sample of 41 sources is substantially smaller than the sample of 96 sources used by Deharvang et al. (2008), despite the larger number of fields used here. This partly reflects our smaller redshift interval ($z = 0.195 - 0.44$), together with our geometric restriction ($R < 32.5$). Only 67 of the Deharvang et al. (2008) sources lie within these bounds: 59 of these are contained in our initial sample of LAEs, while the remaining 8 are marginal lines that fell out of our selection in both the automatic and the visual search. Of the 59 sources, we classified 12 as AGNs based on the UV spectra (mostly on the basis of the line width rather than on the presence of high-excitation lines), and of the remaining 47 sources, only 28 have $\text{EW}(\text{Ly}\alpha) > 20 \text{ Å}$. Thus, 28 of our low-redshift LAE sample overlaps with Deharvang et al. (2008)’s sample, and the remaining sources are either drawn from the other three fields not used in their analysis or are additional sources in the fields in common that were included in our sample but not in their sample. All 28 overlapping sources agree in the *GALEX* redshift, but the Deharvang et al. (2008) $\text{Ly}\alpha$ luminosities are about 10% higher than ours. This may follow from the slight differences in methodology; in particular, we have calculated the fluxes directly from the spectra rather than recalibrating to the broadband fluxes. Thus, 10% is probably a reasonable measure of the systematic uncertainty in the luminosities.

We note that one of the sources in our moderate-redshift LAE sample (GALEX1437+3541) is included in the Deharvang et al. (2008) sample at a redshift of $z = 0.468$. The present redshift of $z = 0.664$ is based on a much stronger emission line seen in the NUV spectrum of this object. The line identified in the FUV by Deharvang et al. is weak and appears to be an artifact. (Note that Deharvang et al. only searched the FUV spectra for $\text{Ly}\alpha$ emission lines.)

5.2. LAE Number Counts




FIG. 13.— Comparison of the number counts as a function of NUV magnitude of the $z = 0.195 - 0.44$ LAEs from our *GALEX* fields with rest-frame $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$ (blue diamonds) and $> 45 \text{ \AA}$ (red triangles) and the number counts of all the NUV continuum sources in our *GALEX* fields (black squares) and in the GOODS-N field (purple solid squares), as well as all the $z = 0.195 - 0.44$ galaxies in the GOODS-N field (purple open squares). The $z = 0.195 - 0.44$ galaxies constitute about 30% of the sources at $\text{NUV} > 21$. The solid curve shows the number counts obtained from the translated $z = 0.2 - 0.4$ NUV-continuum LF of Arnouts et al. (2005). It matches the $z = 0.195 - 0.44$ GOODS-N points reasonably well. The dashed curve shows 5% of the Arnouts et al. (2005) counts, which matches the $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$ LAE counts, and the dotted curve shows 1% of the Arnouts et al. (2005) counts, which matches the $\text{EW}(\text{Ly}\alpha) > 45 \text{ \AA}$ LAE counts.

In Figure 13 we show the number counts per unit magnitude of the LAEs in the redshift interval $z = 0.195 - 0.44$ versus NUV magnitude. This is simply the sum of the inverse areas over all the sources in the magnitude and redshift interval divided by the magnitude interval. We have divided this into two rest-frame $\text{EW}(\text{Ly}\alpha)$ intervals: $> 20 \text{ \AA}$ (blue diamonds) and $> 45 \text{ \AA}$ (red triangles) to show that the EW distribution does not change rapidly with NUV magnitude. We compare the number counts of the LAEs with the number counts of all the NUV continuum sources in our *GALEX* fields (black squares) and in the GOODS-N field (purple solid squares). These points include stars, galaxies, and AGNs. The purple open squares show the number counts of all the $z = 0.195 - 0.44$ galaxies in the GOODS-N sample using the Barger et al. (2008) redshift information and after removing all sources classified as AGNs based on having X-ray luminosities $> 10^{42} \text{ erg s}^{-1}$.

Arnouts et al. (2005) produced a $z = 0.2 - 0.4$ rest-frame $1500 \text{ \AA}$ luminosity function (LF) from *GALEX* VIMOS-VLT Deep Survey data, which we translated to number counts in the same redshift interval versus what corresponds to approximately observed-frame $1900 \text{ \AA}$. We then applied a small -0.2 mag differential K -correction to move that to an NUV magnitude. The result is the solid curve, which agrees reasonably well with the $z = 0.195 - 0.44$ GOODS-N points.

Both the $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$ (blue diamonds) and $\text{EW}(\text{Ly}\alpha) > 45 \text{ \AA}$ (red triangles) LAE number counts show a substantial rise to $\text{NUV} = 21$ where, as we have discussed previously (e.g., Figure 4(a)), the LAEs appear to onset in substantial numbers in the redshift interval

$z = 0.195 - 0.44$. As can be seen from Figure 13, this trend mirrors the overall number counts in the redshift interval $z = 0.195 - 0.44$, which enter the population in significant numbers at magnitudes fainter than 21. We have scaled the Arnouts et al. (2005) curve by 0.05 (dashed) and 0.01 (dotted) to show the fraction of LAEs in a particular $\text{EW}(\text{Ly}\alpha)$ range. Thus, $\sim 5\%$ of the NUV-continuum selected galaxies in this low-redshift interval are LAEs. The curves provide a reasonable approximation to the shape of the LAE counts in both $\text{EW}(\text{Ly}\alpha)$ ranges, showing that the $\text{EW}(\text{Ly}\alpha)$ distribution is not changing rapidly over the observed magnitude range.

5.3. LAE Luminosity Function

We may quantify this further by computing the LAE LF. As Deharveng et al. (2008) stress in their derivation of the LAE LF, the procedure is complicated because we are using a NUV-continuum selected sample and the line flux limits depend on the $\text{EW}(\text{Ly}\alpha)$ distribution. This issue is somewhat alleviated by the relatively invariant rest-frame UV colors above the Lyman continuum break (see the low-redshift sources in Figure 11) so that the NUV limiting magnitude corresponds to an approximate $\text{Ly}\alpha$ flux for a given observed $\text{EW}(\text{Ly}\alpha)$. In principle, by choosing a high enough flux limit in each field we could construct a roughly complete flux-limited sample. This would require us to set the flux limit to correspond to a source with the maximum observed $\text{EW}(\text{Ly}\alpha)$ (about $100 \text{ \AA}$; e.g., Figure 3) at the magnitude limit of the particular field. However, in practice, the present sample is too small to allow such a procedure. An alternative procedure, following Deharveng et al. (2008), is to include all the sources above a lower flux limit than this. We can then attempt to correct for the incompleteness by assuming the $\text{EW}(\text{Ly}\alpha)$ distribution of Figure 3 is invariant as a function of NUV magnitude. This allows us to correct for the missing high- $\text{EW}(\text{Ly}\alpha)$ sources with NUV magnitudes fainter than the magnitude limit of the field. This assumption of invariance in the $\text{EW}(\text{Ly}\alpha)$ distribution may well fail as we move to fainter magnitudes, where we may see higher $\text{EW}(\text{Ly}\alpha)$ sources, so it is important to minimize the extrapolation. We use the latter procedure here.

We first set the $\text{Ly}\alpha$ flux limits high enough—corresponding to a rest-frame $\text{EW}(\text{Ly}\alpha) > 45 \text{ \AA}$ at the limiting NUV magnitude of each field—both to include a significant fraction of the sources in Figure 3 and to minimize the incompleteness corrections. The downside of this high flux cut is that we reduce the already small sample significantly and are restricted to high $\text{Ly}\alpha$ luminosities. However, the incompleteness corrections are small. We also try a low flux cut with a flux limit corresponding to a rest-frame $\text{EW}(\text{Ly}\alpha) > 25 \text{ \AA}$ at the limiting NUV magnitude of each field, which allows us to probe to lower luminosities at the expense of a higher incompleteness correction.

Specifically, for each field we determined a $\text{Ly}\alpha$ flux limit corresponding to a source with the limiting NUV magnitude of Table 1 corrected to an FUV magnitude by adding an offset of 0.37 mag (see Figure 11) and with an observed $\text{EW}(\text{Ly}\alpha)$ corresponding to the chosen rest-frame limit placed at the center of the redshift interval. For a rest-frame $\text{EW}(\text{Ly}\alpha)$ of $45 \text{ \AA}$, the adopted flux limit




FIG. 14.— LAE (rest-frame $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$) LF in the redshift interval $z = 0.195 - 0.44$ using two flux limits (see text). The red diamonds show the LF using the high flux cut where incompleteness corrections are minimal, and the black squares show the LF using the low flux cut that extends to lower luminosities, but where the incompleteness corrections are larger. In each case the raw (incompleteness-corrected) LAE LF is shown with open (solid) symbols. The error bars are $\pm 1\sigma$ based on the Poisson errors for the number of sources in each luminosity bin. The solid curve shows the LAE LF obtained by convolving the $\text{EW}(\text{Ly}\alpha)$ distribution of Figure 3 with the translated $z = 0.2 - 0.4$ Arnouts et al. (2005) NUV-continuum LF. The dashed curve shows the same but using the $\text{EW}(\text{Ly}\alpha)$ distribution of the $\text{NUV} < 21.8$ galaxies only, which should roughly match to the raw LAE LFs. The blue triangles show Deharvang et al. (2008)’s determination of the LAE LF before (open) and after (solid) their incompleteness correction. We have reduced Deharvang et al. (2008)’s luminosity scale by 10% to make the $\text{Ly}\alpha$ luminosity measurements consistent (see Section 5.1).

is $f(\text{Ly}\alpha) = 3.4 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ for the GROTH 00 field; it is proportionally higher for the remaining shallower fields. Only galaxies with fluxes above each field’s adopted flux limit were included in the final sample. The observed area at a given flux is then the sum of all the field areas where the limiting flux is lower than this flux.

We next constructed the LAE LF in the redshift interval $z = 0.195 - 0.44$ using the $1/V$ technique (Felten 1976). In Figure 14 we use red open diamonds (black open squares) to show the LF for the high flux (low flux) cut. The $\pm 1\sigma$ errors are based on the Poisson errors corresponding to the number of sources in each bin. Since the line width split that we used to separate out AGNs is somewhat subjective, as a check we have also calculated the LF including all the sources in the redshift interval that do not have high-excitation lines. This slightly increases the LF but by an amount which is small compared to the uncertainties.

These are the raw LFs, without the incompleteness corrections for the missing high- $\text{EW}(\text{Ly}\alpha)$ sources in the flux-limited samples. In order to compute the incompleteness corrections, we used the form of the Arnouts et al. (2005) LF to obtain the expected number of continuum sources in the redshift interval at fainter UV magnitudes. We then drew the correct number of sources from the $\text{EW}(\text{Ly}\alpha)$ distribution of Figure 3 to simulate the missing high- $\text{EW}(\text{Ly}\alpha)$ sources and recomputed the LFs. The incompleteness-corrected LFs are shown with the solid symbols in Figure 14. In general these correc-

tions are small, except in the faintest bin of the low flux cut sample (black squares).

Given our assumption of an invariant $\text{EW}(\text{Ly}\alpha)$ distribution, it is also possible to simply convolve this distribution with the observed NUV-continuum LF in the redshift interval to determine the LAE LF. This allows us to construct a LF to fainter luminosities but at the expense of assuming the same $\text{EW}(\text{Ly}\alpha)$ distribution applies at substantially fainter NUV magnitudes than where it was measured. As we have stressed above, the invariant $\text{EW}(\text{Ly}\alpha)$ distribution assumption may not be valid if fainter sources have different $\text{Ly}\alpha$ emission-line properties. We did the calculation using the $\text{EW}(\text{Ly}\alpha)$ distribution of Figure 3 and the translated $z = 0.2 - 0.4$ Arnouts et al. (2005) NUV-continuum LF. This result is shown with the solid curve in Figure 14. We repeated the calculation using the $\text{EW}(\text{Ly}\alpha)$ distribution of the $\text{NUV} < 21.8$ galaxies only to illustrate the luminosity range corresponding to the actual measurements. This result is shown with the dashed curve. The incompleteness corrections can then be measured from the ratio of the two curves (this more closely mirrors the procedure used by Deharvang et al. 2008), and they agree well with what we found previously.

All of our LAE LF measurements are comparable to the raw LAE LF determined by Deharvang et al. (2008; blue open triangles in Figure 14). However, we do not find the large incompleteness corrections that they found (the blue solid triangles show their incompleteness-corrected LAE LF). Their corrections appear remarkably large, particularly at the high-luminosity end. There may be differences reflecting the selection in the $\text{EW}(\text{Ly}\alpha)$ (here we are using a rigid rest-frame $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$ definition of the LAEs) and the more stringent exclusion of potential AGNs in the present analysis, but it does not appear that these can account for the differences. The problem may be caused by a missing color correction in the Deharvang analysis (J.-M. Deharvang, priv. comm.).

In Figure 15 we compare our $z = 0.195 - 0.44$ LAE LF (here we adopt the low flux cut incompleteness-corrected result; black squares), which was chosen with the same rest-frame $\text{EW}(\text{Ly}\alpha)$ selection (i.e., $> 20 \text{ \AA}$) as the high-redshift samples, with the $z = 3.1$ LAE LF of Gronwall et al. (2007; red curve). Other determinations of the $z \sim 3$ LAE LF are extremely similar (e.g., Cowie & Hu 1998; van Breukelen et al. 2005; Ouchi et al. 2008). We have made a maximum likelihood analysis of our data to obtain a Schechter (1976) function fit. Given the limited dynamic range of our data, we have held the slope α fixed and only measured L_* . For the Gronwall et al. (2007) slope of $\alpha = -1.36$, we obtain $\log L_* = 41.76 \pm 0.08$. Normalizing to the observed number of sources gives $\phi_* = (1.95 \pm 0.35) \times 10^{-4} \text{ Mpc}^{-3}$. The errors are $\pm 1\sigma$. This fit is shown as the blue curve in Figure 15. For the van Breukelen et al. (2005) slope of $\alpha = -1.6$, we obtain $\log L_* = 41.81 \pm 0.09$. Normalizing to the observed number of sources gives $\phi_* = (1.68 \pm 0.30) \times 10^{-4} \text{ Mpc}^{-3}$.

Using the Gronwall et al. (2007) slope and comparing to the Gronwall et al. (2007) LF gives a drop of ~ 8 in L_* and ~ 6.5 in ϕ_* for a drop in the $\text{Ly}\alpha$ luminosity density of ~ 50 . Using the van Breukelen et al. (2005) LF gives a




FIG. 15.— Comparison of our $z = 0.195 - 0.44$ low flux cut incompleteness-corrected LAE LF (black squares) with the $z = 3.1$ LAE LF of Gronwall et al. (2007; red curve). Other determinations of the $z \sim 3$ LAE LF are extremely similar (see text). The blue curve shows the maximum likelihood Schechter (1976) function fit to our data for a fixed slope of $\alpha = -1.36$ (that of Gronwall et al. 2007). There is a drop of ~ 8 in L_* and ~ 6.5 in the normalization ϕ_* relative to the $z = 3.1$ LAE LF. The dotted black curve shows the LAE LF obtained by convolving the $\text{EW}(\text{Ly}\alpha)$ distribution of Figure 3 with the translated $z = 0.2 - 0.4$ Arnouts et al. (2005) NUV-continuum LF.

drop in the $\text{Ly}\alpha$ luminosity density of 55.

As Deharveng et al. (2008) have pointed out, this substantial drop in the LF from higher redshift values is considerably in excess of the corresponding continuum UV light density drop. Thus, it appears that LAEs are far less common now than they were in the past and that they have lower luminosities.

5.4. LAE Equivalent Widths

In order to compare the LAE $\text{EW}(\text{Ly}\alpha)$ distributions at low and high redshifts, we need to translate our measured $\text{EW}(\text{Ly}\alpha)$ distribution as a function of NUV magnitude (Figure 3) into one which is a function of LAE luminosity. To do this we computed the number density of LAEs in the redshift interval $z = 0.195 - 0.44$ as a function of their rest-frame $\text{EW}(\text{Ly}\alpha)$ using the same methodology that we used to compute the incompleteness-corrected LAE LFs. We used a low flux cut corresponding to a rest-frame $\text{EW}(\text{Ly}\alpha)$ of $25 \text{ \AA}$ to probe to low luminosities and computed the number density of sources above a limiting luminosity of $4 \times 10^{41} \text{ erg s}^{-1}$ by summing the inverse volumes of all the sources in the $\text{EW}(\text{Ly}\alpha)$ interval lying above this luminosity. We then divided by the width of the $\text{EW}(\text{Ly}\alpha)$ bin. The results are shown in Figure 16.

The low-redshift LAE $\text{EW}(\text{Ly}\alpha)$ distribution is well described by the same exponential with a scale length of $75 \text{ \AA}$ that provides a good fit to the $z \sim 3$ LAE $\text{EW}(\text{Ly}\alpha)$ distribution (Gronwall et al. 2007). This is shown by the red solid curve in Figure 16. Thus, the form of the $\text{EW}(\text{Ly}\alpha)$ distribution for the LAEs is not changing with redshift, even though the number of sources satisfying the LAE criterion is much smaller at low redshifts. We do not see any extreme EWs($\text{Ly}\alpha$) (greater than $120 \text{ \AA}$) in the present sample, but this may be a




FIG. 16.— Number density of sources above $\log L(\text{Ly}\alpha) = 41.6 \text{ erg s}^{-1}$ vs. rest-frame $\text{EW}(\text{Ly}\alpha)$. The number densities correspond to a $10 \text{ \AA}$ bin size. The red solid curve shows an exponential fit to the data with a $75 \text{ \AA}$ scale length equal to that in the $z = 3.1$ population, which provides a good description of the data. The blue dashed curve shows the steeper fit to the continuum selected data of Figure 3.

simple consequence of the continuum selection, which is biased against finding such objects.

5.5. Ly-alpha Velocities Relative to H-alpha

A comparison of the redshifts of the $\text{Ly}\alpha$ lines relative to the redshifts of the $\text{H}\alpha$ lines is of considerable interest since it may relate to the kinematical structure of the galaxy and the escape process of the $\text{Ly}\alpha$ photons. However, this comparison is difficult because of the wavelength calibration uncertainties in the *GALEX* grism data. Morrissey et al. (2007) give a calibration error for the *GALEX* wavelengths of about 3% in the body of each spectral order and about 10% near the edges of each order.

In Figure 17 we show the wavelength offsets between the $\text{Ly}\alpha$ wavelength that would be measured from *GALEX* and the $\text{Ly}\alpha$ wavelength that would be inferred from the optical redshifts. For this figure we have augmented the *GALEX* LAE sample (black solid squares; this includes any source where we have measured the redshift from our optical data, as well as any source where the optical redshift could be obtained from the NED database⁷) in order to obtain a larger number of sources to maximize our understanding of the effect. The additional sources are AGNs (red solid diamonds are based on the presence of high-excitation lines; red open squares are based on the line widths) in our *GALEX* fields with optical redshifts, again either from our own observations or from NED.

Nearly all of the sources show a *GALEX* $\text{Ly}\alpha$ wavelength that is redder than would be expected from the optical redshift. The median offset is $3.9 \text{ \AA}$ for the sources (both AGNs and galaxies) in the redshift interval $z = 0.195 - 0.44$. Given that both the AGNs and the

⁷ The NASA/IPAC Extragalactic Database (NED) is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration




FIG. 17.— The offset of the observed Ly α wavelength in the *GALEX* spectra from that expected from the optical redshift vs. wavelength. Black solid squares show the LAEs, red solid diamonds show AGNs with high-excitation lines in the UV spectra, and red open squares show sources without high-excitation lines but with line widths greater than 15 Å. The optical redshifts come from our observations and from NED.

LAEs show this offset, it is probable that much of it is indeed due to the absolute calibration uncertainty. The median wavelength offset is 3.9 Å for just the LAEs and 3.8 Å for just the AGNs. The median redshifting seen in $z \sim 2$ galaxies, which is around 550 km s $^{-1}$ (Pettini et al. 2001), would produce an offset of about 2.2 Å, which is well within the uncertainties.

6. PROPERTIES OF THE LOW-REDSHIFT LY α GALAXY SAMPLE

6.1. Ly α Versus H α Fluxes

Focusing on our optical spectroscopic observations of the *GALEX* spectroscopic sample, we now turn to analyzing the properties of the optically-confirmed Ly α Galaxies in the redshift interval $z = 0.195 - 0.44$ (Table 11, excluding the sources classified as AGNs). We compare them to the properties of both the *GALEX* spectroscopic sample without UV spectral identifications that are optically classified as galaxies and lie in the same redshift interval (Table 13; hereafter, we refer to this as the optically-confirmed NUV-continuum selected galaxy sample) and the essentially completely spectroscopically identified GOODS-N NUV < 24 galaxy sample in the redshift interval $z = 0.15 - 0.48$ (hereafter, we refer to this as the GOODS-N NUV-continuum selected galaxy sample). In the latter sample we have removed all the sources classified as AGNs based on having X-ray luminosities > 10 42 erg s $^{-1}$. There are no sources in the GOODS-N sample that appear in the *GALEX* Ly α Galaxy sample. However, we caution that there may still be some Ly α emission-line galaxies in the GOODS-N sample, since the *GALEX* grism data for this field do not go as faint in NUV as the optical spectroscopy.

In Figure 18 we show the ratio of the Ly α flux to the H α flux versus Ly α luminosity. The optically-confirmed Ly α Galaxies with rest-frame EW(Ly α) > 20 Å (red solid triangles) mostly lie in a fairly narrow region from just below 1 to a maximum ratio of 8. The median value

of 2.2 ± 1.0 is shown by the red solid line, while the average ratio is 2.6. This value is about 4 times smaller than the case B ratio (which, depending on the electron density, is 8 – 12; Ferland & Osterbrock 1985) often used to translate Ly α luminosity to star formation rate, and it is consistent with the similar reduction inferred by comparing star formation rates measured from the UV continuum with those measured from Ly α in the $z \sim 3$ emitters (Gronwall et al. 2007). It is also consistent with the range of values seen in other optical spectroscopic follow-up observations of the Deharvang et al. (2008) sample (Atek et al. 2009; Mallery 2009). The galaxies with detected Ly α lines but rest-frame EW(Ly α) weaker than 20 Å are shown with green very small solid triangles. These generally have lower ratios of Ly α /H α , with a median value of 1.03, though the number of objects is too small to derive a median error. There appears to be no obvious dependence on other parameters such as metallicity though a large sample is clearly required to explore this in depth.

If we use Ly α /H α = 2.6 and adopt the widely used Kennicutt (1998) conversion of the H α luminosity to star formation rate for the Salpeter (1955) initial mass function (IMF) extending to 0.1 M $_{\odot}$, we obtain a conversion of the Ly α luminosity to star formation rate of

$$\log \text{SFR} = -40.67 + \log L(\text{Ly}\alpha). \quad (1)$$

The true value for the H α calibration depends on the time history of the star formation. The value derived in Cowie & Barger (2008), which is appropriate for the average of galaxies at these redshifts, would reduce this by 0.2 dex to $-40.47 + \log L(\text{Ly}\alpha)$.

While the above equation can be used to get the star formation rates for Ly α galaxies selected as in this sample—a selection which corresponds fairly closely to the formal LAE definition—it must be noted that the observed values of Ly α /H α in this sample lie above a threshold set by the EW(Ly α) selection (see Figure 3). In other words, if the EW(Ly α) is too small, we will not see a Ly α line. Thus, the narrow range of observed values in Figure 18 is merely a selection bias, as we illustrate by putting upper bounds on the optically-confirmed NUV-continuum selected galaxies (blue diamonds). These stretch up to overlap the lower boundary of the optically-confirmed Ly α Galaxy sample.

The reason for the selection bias is as follows. A given UV continuum luminosity corresponds to an approximate H α luminosity in the absence of continuum extinction, and for a given EW(Ly α), it also specifies the Ly α luminosity. For this sample we can use the empirical star formation rate calibrations given in Cowie & Barger (2008),

$$\log \text{SFR} = -42.63 + \log L(\text{FUV}), \quad (2)$$

$$\log \text{SFR} = -40.90 + \log L(\text{H}\alpha), \quad (3)$$

and the conversion from FUV fluxes to NUV fluxes based on Figure 11 (NUV=FUV−0.37 mag) to roughly obtain a relation between the NUV continuum luminosity and the H α luminosity. Then we use the NUV continuum flux and the EW(Ly α) to determine the Ly α luminosity and obtain the relation

$$L(\text{Ly}\alpha)/L(\text{H}\alpha) = 2.4(\text{EW}(\text{Ly}\alpha)/20 \text{ Å}), \quad (4)$$

where EW(Ly α) is the rest-frame EW(Ly α). Thus, the




FIG. 18.— $\text{Ly}\alpha/\text{H}\alpha$ flux ratio vs. $\text{Ly}\alpha$ luminosity for the *GALEX* sources that appear to be star formers based on their UV and optical spectra, lie in the redshift interval $z = 0.195 - 0.44$, and have SDSS magnitudes. The red small solid triangles show the measured ratios for the optically-confirmed $\text{Ly}\alpha$ Galaxies with rest-frame $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$. The red solid line shows the median ratio of 2.1 for this sample. The red large solid triangles show the sources with $\text{EW}(\text{Ly}\alpha) > 40 \text{ \AA}$, and the green very small solid triangles show the sources with detected $\text{Ly}\alpha$ and $\text{EW}(\text{Ly}\alpha) \leq 20 \text{ \AA}$. All error bars are $\pm 1\sigma$. The red dotted line marks the lower limit of the ratios ($8 - 12$) expected for case B (Ferland & Osterbrock 1985). The blue open diamonds with the downward pointing arrows denote the upper limits on the $\text{Ly}\alpha/\text{H}\alpha$ flux ratios for the optically-confirmed NUV-continuum selected galaxies. Here we have assumed an observed-frame $\text{EW}(\text{Ly}\alpha) = 10 \text{ \AA}$ in calculating the upper limits for the flux ratios and the $\text{Ly}\alpha$ luminosities.

effect of the $\text{EW}(\text{Ly}\alpha)$ selection is to place a lower bound on $L(\text{Ly}\alpha)/L(\text{H}\alpha)$, while case B places an upper bound. The exact lower bound does depend on the details of the star formation history, which determines the exact UV continuum to $\text{H}\alpha$ conversion, and on the extinction, which can lower the $L(\text{Ly}\alpha)/L(\text{H}\alpha)$ ratios. However, the lower limit bias will still be present. Thus, we cannot simply estimate general escape fractions from $\text{Ly}\alpha$ galaxy samples alone, because we also need to deal with all of the $\text{Ly}\alpha$ undetected sources, which still produce some $\text{Ly}\alpha$ emission.

With larger samples we could begin to determine the incompleteness corrections as a function of EW. Here we follow a simpler route of using a stacking analysis to determine the $L(\text{Ly}\alpha)/L(\text{H}\alpha)$ ratio in the sources from the NUV-continuum sample that lie in the redshift range but do not have identified $\text{Ly}\alpha$. We can then estimate the fraction of $\text{Ly}\alpha$ photons that escape from NUV-continuum selected galaxies at a particular redshift using the ratio of the total $\text{Ly}\alpha$ light to the total $\text{H}\alpha$ light and combine this with the measured values in the objects with identified $\text{Ly}\alpha$ to form an average appropriate for the total galaxy sample in the redshift range.

To do this, we constructed average rest-frame galaxy spectra. These are shown in Figure 19 for three samples: rest-frame $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$ (red spectrum), detected $\text{Ly}\alpha$ galaxies with $\text{EW}(\text{Ly}\alpha) \leq 20 \text{ \AA}$ (green spectrum), and NUV-continuum selected objects (blue spectrum). The averaged spectra are constructed using the optically measured redshifts with an average offset deter-




FIG. 19.— Average *GALEX* spectrum for the objects with rest-frame $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$ (red spectrum), detected $\text{Ly}\alpha$ with rest-frame $\text{EW}(\text{Ly}\alpha) \leq 20 \text{ \AA}$ (green spectrum), and NUV-continuum selected objects (blue spectrum). The vertical dotted lines show the strong features in the spectra. In (a) we show the redshift interval $0.195 - 0.28$, and in (b) we show the redshift interval $0.28 - 0.44$.

mined from Figure 17 to match the *GALEX* wavelength calibration. The strong metal absorption lines produce weak features in the low-resolution spectra, which are marked on the figure with vertical dotted lines. To simplify the averaging procedure, we have divided the redshift range into two intervals: (a) $z = 0.195 - 0.28$ and (b) $z = 0.28 - 0.44$. This split has the secondary advantage of allowing us to see that the results are robust and reproducible with different samples.

We can see from Figure 19 that the NUV-continuum selected galaxies do have weak $\text{Ly}\alpha$ emission. We measured the $\text{Ly}\alpha$ flux and compared it to the average $\text{H}\alpha$ flux of the same sample to determine an average $\text{Ly}\alpha/\text{H}\alpha$ ratio for these sources. For the full $0.195 - 0.44$ range we find a value of 0.23, but both redshift ranges shown in Figure 19 give similar values. However, for these weaker sources there may be a substantial correction for the presence of $\text{Ly}\alpha$ absorption in some of the galaxies. A maximum estimate for this can be obtained by measuring the line flux relative to the zero level rather than relative to the continuum. This gives a maximum $\text{Ly}\alpha/\text{H}\alpha$ ratio for the sources of 0.54.




FIG. 20.— (a) Comparison of the $z = 0.195 - 0.44$ H α (red squares), H β (blue diamonds), and [OII] $\lambda 3727$ (cyan triangles) emission-line selected galaxy LFs from the GOODS-N data with the LAE LF (black squares and black dotted curve). The red solid curve shows the Tresse & Maddox (1998) Schechter (1976) function fit to their H α LF at $z = 0.2$ scaled to the present geometry. (b) As in (a), but the luminosities of the lines have now been scaled to the H α luminosity. The black dashed curve shows the factor of 20 change in normalization required to bring the LAE LF into rough consistency with the other emission-line LFs.

If we make a weighted addition of the 5% of LAEs with rest-frame $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$ and $\text{Ly}\alpha/\text{H}\alpha$ ratios of 2.6, the 2% of LAEs with $\text{EW}(\text{Ly}\alpha) \leq 20 \text{ \AA}$ and $\text{Ly}\alpha/\text{H}\alpha$ ratios of 1.0, and the 93% of non-LAEs with $\text{Ly}\alpha/\text{H}\alpha$ ratios of 0.23 or 0.54, we obtain an average $\text{Ly}\alpha/\text{H}\alpha$ ratio of 0.36 – 0.65 at this redshift. Comparing this with the case B ratio, we find that, on average, about 3 – 8% of the Ly α photons are escaping from the entire galaxy population at $z = 0.3$. A minimum of a quarter of these photons are emerging from the small fraction of the identified LAEs alone, and the fraction could be as high as 40%.

We may also look at the issue of the overall Ly α escape fraction by comparing the LFs of the LAEs with the LFs of other emission-line selected galaxies. For the redshift interval $z = 0.195 - 0.44$ we show in Figure 20(a) the LAE LF (black squares and dotted curve) compared with the H α (red squares), H β (blue diamonds), and [OII] $\lambda 3727$ (cyan triangles) emission-line selected galaxy LFs from the GOODS-N sample of Barger et al. (2008).




FIG. 21.— Rest-frame $\text{EW}(\text{Ly}\alpha)$ vs. rest-frame $\text{EW}(\text{H}\alpha)$ for the GALEX sources that appear to be star formers based on their optical spectra and that lie in the redshift interval $z = 0.195 - 0.44$. These include the optically-confirmed Ly α Galaxies (red solid triangles) from Table 12 and the optically-measured NUV-continuum selected galaxies (blue open diamonds at a nominal value of 6 Å) from Table 14. The $\pm 1\sigma$ error bars are shown for the $\text{EW}(\text{Ly}\alpha)$ of the Ly α Galaxies. In many cases the statistical error in $\text{EW}(\text{H}\alpha)$ is too small to be seen, and we have instead shown a 10% systematic error. The black vertical dashed line shows Kakazu et al. (2007)'s definition of ultra-strong emission-line galaxies. This roughly matches the $\text{EW}(\text{H}\alpha)$ above which a very large fraction of the sources are Ly α Galaxies.

We also show the $z = 0.2$ H α LF of Tresse & Maddox (1998) rescaled to the present geometry (red curve), which agrees well with the GOODS-N H α LF. In Figure 20(b) we show the LFs for each of the emission lines rescaled to the H α luminosity. For [OII] and H β we made the conversion using the star formation rate relations relative to H α from Cowie & Barger (2008), while for Ly α we made the conversion with the ratio of 2.6 derived above.

We can see from Figure 20(b) that the other emission-line LFs give fully consistent LFs when placed on the common scale. However, the LAE LF, while similar in shape, is much lower in normalization. This again emphasizes that the LAEs comprise only a fraction of the star-forming galaxies at these redshifts. The LAE LF can be renormalized to match the H α LF by multiplying by a factor of roughly 20. This is shown as the dashed black curve in Figure 20(b). Thus, this alternative analysis also suggests that 5% of the Ly α light is escaping at this redshift.

6.2. Ly α versus H α Equivalent Widths

One of the most interesting questions is whether there is a way to pick out Ly α emission-line galaxies using only optical spectra. The best diagnostic seems to be the magnitude of the EW in the Balmer lines. In Figure 21 we show the $\text{EW}(\text{Ly}\alpha)$ versus the $\text{EW}(\text{H}\alpha)$ for the optically-confirmed Ly α Galaxies (red solid triangles) and for the optically-confirmed NUV-continuum selected galaxies (blue open diamonds). Overall, the $\text{EW}(\text{H}\alpha)$ for the Ly α Galaxies (median value of 76 Å) are significantly higher than those for the NUV-continuum selected galaxies (median value of 36 Å). Roughly half have a rest-

frame $\text{EW}(\text{H}\alpha) > 80 \text{ \AA}$. Kakazu et al. (2007) call such sources ultra-strong emission-line galaxies (USELs). In our optically-identified sample, we see that all but two of the USELs are $\text{Ly}\alpha$ Galaxies and all of the very high $\text{EW}(\text{H}\alpha)$ sources are. Clearly we have optically observed a much smaller fraction of the NUV-continuum selected sample than of the candidate $\text{Ly}\alpha$ Galaxy sample, so it is possible that we might see more scattering into the very high $\text{EW}(\text{H}\alpha)$ region with more observations. However, the current data strongly suggest that a large fraction of USELs are $\text{Ly}\alpha$ Galaxies.

As is well known, the $\text{EW}(\text{H}\alpha)$ can give a rough estimate of the age of the star formation in a galaxy. For a Salpeter (1955) IMF and a constant star formation rate, the $\text{EW}(\text{H}\alpha)$ would drop smoothly to a value of $80 \text{ \AA}$ at about 10^9 yr (Leitherer et al. 1999), while an instantaneous starburst would drop below this value after about 10^7 yr . It is therefore possible that the presence of a high $\text{EW}(\text{Ly}\alpha)$ is simply an age effect, with the youngest galaxies having the strongest $\text{Ly}\alpha$ emission. However, it could also be that there are other effects that let the $\text{Ly}\alpha$ photons out more easily, such as $\text{Ly}\alpha$ galaxies having lower metallicities or more kinematic disturbances than the general population. We now turn to the measurement of these quantities.

6.3. Metallicities

Only a small number of the sources have a detectable $[\text{OIII}]\lambda 4363$ line that we can use to make a direct estimate of the O abundance. We will discuss these at the end of the section. For our primary analysis of the metallicities we use the $\text{N2} = \log([\text{NII}]\lambda 6584/\text{H}\alpha)$ diagnostic ratio (Storchi-Bergmann et al. 1994). We use N2 since the $\text{R23} = (1.3[\text{OIII}]\lambda 5007 + [\text{OII}]\lambda 3727)/\text{H}\beta$ diagnostic ratio of Pagel et al. (1979) is multivalued over the metallicity range of interest (McGaugh 1991), and our spectral flux calibration is not adequate to use the $[\text{NII}]/[\text{OII}]$ ratio. The N2 diagnostic also has the advantage of being the method used by Erb et al. (2006) to estimate the metallicities of their $z \sim 2$ galaxy sample. The downside of the N2 diagnostic is that it is highly sensitive to the ionization parameter (e.g., Kewley & Dopita 2002). Other drawbacks to the N2 diagnostic include variations in N/O, and its sensitivity to contamination by a high- $[\text{NII}]/\text{H}\alpha$ AGN contribution.

Pettini & Pagel (2004) showed that locally there is a reasonably tight relation between $12 + \log(\text{O}/\text{H})$ and N2 for systems where the O abundance has been determined with the direct method; their linear fit gives $12 + \log(\text{O}/\text{H}) = 8.90 + 0.57\text{N2}$ over the range $\text{N2} = -2.5$ to -0.5 . Extrapolating this to higher redshifts requires assuming that there is no change in the typical ionization parameter, which may well be incorrect. However, at $z = 0.195 - 0.44$ Cowie & Barger (2008) found a narrow range of ionization parameters ($q \sim 2 \times 10^7$) that, when combined with the photoionization code-based estimates of Kobulnicky & Kewley (2004), gives a broadly similar though considerably more analytically complex equation to the Pettini & Pagel (2004) relation. Cowie & Barger (2008) also showed that other line diagnostics gave similar metallicity-luminosity relations to that derived from N2 in the redshift interval $z = 0.195 - 0.44$. The maximum deviation between the relation used in Cowie &

Barger (2008) and the Pettini & Pagel (2004) linear relation over the range -2 to -0.5 is -0.26 dex at -2 and $+0.26 \text{ dex}$ at -0.5 . The local data may be slightly better represented by the Cowie & Barger (2008) relation over this range, though the differences are probably not very meaningful. In the following we will use the Pettini & Pagel (2004) linear relation for simplicity and to allow a direct comparison with the high-redshift results, but we will always show the measured N2 as our primary variable.

In Figure 22(a) we show the Baldwin et al. (1981; BPT) diagnostic diagram of $[\text{OIII}]\lambda 5007/\text{H}\beta$ versus $[\text{NII}]\lambda 6584/\text{H}\alpha$ for the *GALEX* sources that are not classified as AGNs based on their UV spectra, do not show broad lines in their optical spectra, and lie in the redshift interval $z = 0.195 - 0.44$. Only spectra with a significantly detected $\text{H}\beta$ line (signal-to-noise greater than three) are included in the diagram. These include the optically-confirmed $\text{Ly}\alpha$ Galaxies (red solid triangles) and the optically-confirmed NUV-continuum selected galaxies (blue open diamonds). We also show the GOODS-N NUV-continuum selected galaxy sample with $\text{NUV} = 20 - 22$ and redshifts $z = 0.15 - 0.48$ (blue solid diamonds). The diagram separates AGN-dominated sources from star-forming galaxies. The dotted curve is the maximum starburst curve of Kewley et al. (2001), and the dashed curve is the Kauffmann et al. (2003) star former/AGN separator from the SDSS data. Sources lying well above these curves are AGNs. Although most of our sources lie along the star-forming galaxy track, the diagram suggests that one of the optically-confirmed $\text{Ly}\alpha$ Galaxies (GALEX1240+6233) is in fact an AGN. We have eliminated this object from all other figures, except the BPT diagrams of Figure 29. Finkelstein et al. (2009) similarly see a small number of BPT-selected AGNs in their analysis of the Deharvang et al. (2008) $\text{Ly}\alpha$ sample.

Finkelstein et al. (2009) also classified GALEX1417+5228 as an AGN based on the presence of $\text{HeII}\lambda 4686$ in the spectrum. Three of the present optical spectra show detectable HeII : GALEX1001+0233 ($f(\text{HeII})/f(\text{H}\beta) = 0.019 \pm 0.005$), GALEX1240+6233 ($f(\text{HeII})/f(\text{H}\beta) = 0.072 \pm 0.015$), and GALEX1417+5228 ($f(\text{HeII})/f(\text{H}\beta) = 0.018 \pm 0.005$). GALEX1240+6233 is classified as an AGN based on the BPT diagram, and the strong HeII confirms this interpretation. However, we do not think that the presence of the weaker HeII lines necessarily implies that the other two objects are AGNs rather than metal-poor star-forming galaxies. Roughly 10% of blue compact galaxies have detectable HeII at the level seen in these galaxies, probably produced by Wolf-Rayet stars or shocks in the galaxies, and the presence of the line becomes more common as one moves to more metal-poor galaxies (e.g., Thuan & Izotov 2005). GALEX1001+0233 and GALEX1417+5228 also have very weak or undetected $\text{NII}\lambda 6584$ which would require them to be metal-poor AGNs. While a very small number of such objects have now been found (Izotov & Thuan 2008), they would not be expected to lie on the low-metallicity star track in the BPT diagram as the present objects do (Groves et al. 2006). In the case of GALEX1001+0233, where the spectrum covers $\text{NeV}\lambda 3426$, there is no sign of this high-excitation




FIG. 22.— (a) BPT diagram for the *GALEX* sources that do not contain broad lines in their optical spectra and that lie in the redshift interval $z = 0.195 - 0.44$. Only objects with significant H β line detections ($> 3\sigma$) and where the spectrum covers the [NII] $\lambda 6584$ line are shown, giving 27 optically-confirmed Ly α Galaxies (red solid triangles) and 19 optically-confirmed NUV-continuum selected galaxies (blue open diamonds). The error bars are $\pm 1\sigma$. We also show the GOODS-N NUV-continuum selected galaxy sample with $NUV = 20 - 22$ and redshifts between $z = 0.15$ and $z = 0.48$ (blue small solid diamonds). The dotted curve is the maximum starburst curve of Kewley et al. (2001), and the dashed curve is the Kauffmann et al. (2003) star former/AGN separator from the SDSS data. The black large open square denotes the source GALEX1240+6233, which would be classified as an AGN on the basis of this diagram. (b) Number of sources vs. $\log [NII]/H\alpha$. All sources with high signal-to-noise ($> 10\sigma$) H α detections and where the [NII] $\lambda 6584$ line is covered by the spectrum are shown. The red shaded histogram shows the 30 optically-confirmed Ly α Galaxies selected in this way, the blue open histogram shows the 29 optically-confirmed NUV-continuum selected galaxies, and the blue shaded histogram (upper part of diagram) shows the GOODS-N NUV-continuum selected galaxies.

line. We therefore classify GALEX1001+0233 and GALEX1417+5228 as star formers.

Sources lying at lower values of [NII] $\lambda 6584/H\alpha$ correspond to lower metallicity galaxies, and we can see immediately from Figure 22(a) that the optically-confirmed

Ly α Galaxies have lower values of [NII] $\lambda 6584/H\alpha$ than the combined NUV-continuum selected galaxies. This may be more clearly seen in Figure 22(b), where we show the distribution of $\log([NII]\lambda 6584/H\alpha)$ for the optically-confirmed Ly α Galaxies (red shaded histogram), for the optically-confirmed NUV-continuum selected galaxies (blue open histogram), and for the GOODS-N NUV-continuum selected galaxies with $NUV = 20 - 22$ (upper blue shaded histogram). While the Ly α Galaxy distribution overlaps with the NUV-continuum selected galaxy distribution, it clearly extends to lower values, and the median [NII] $\lambda 6584/H\alpha$ is lower. A rank sum test gives a 4×10^{-4} probability that the two *GALEX* samples are similar and only a 9×10^{-5} probability that the Ly α Galaxy sample is drawn from the combined NUV-continuum selected samples from both *GALEX* and the GOODS-N.

Given the metallicity-luminosity relation, where lower luminosity sources also have lower metallicities, this result could mean that the Ly α Galaxies are simply lower luminosity galaxies than the NUV-continuum selected galaxies. In order to test this we show in Figure 23 [NII] $\lambda 6584/H\alpha$ versus absolute rest-frame B magnitude, $M_B(AB)$, for the optically-confirmed Ly α Galaxies (red solid triangles) and for the optically-confirmed NUV-continuum selected galaxies (blue open diamonds). (We have not attempted to construct the corresponding metallicity-mass relation because of the difficulty of computing the mass from the optical magnitudes in these star formation dominated objects with strong emission lines.) The values of $12 + \log(O/H)$ computed from the Pettini & Pagel (2004) linear relation are shown on the right-hand axis of the figure. Only galaxies with SDSS magnitudes are shown, and we interpolated between the SDSS model C g and r magnitudes to obtain $M_B(AB)$. We also show in the figure the GOODS-N NUV-continuum selected galaxy sample with $NUV < 24$ and redshifts in the interval $z = 0.15 - 0.48$ (blue solid diamonds). Here the $M_B(AB)$ magnitudes are computed from the AUTO magnitudes in the F606W bandpass of the ACS GOODS-N data (Giavalisco et al. 2004). As would be expected if the photometry is consistent, the *GALEX* optically-confirmed NUV-continuum selected galaxies (blue open diamonds) match to the bright end of the GOODS-N NUV-continuum selected galaxies (blue solid diamonds). The optically-confirmed Ly α Galaxies (red solid triangles) lie systematically lower in metallicity at a given luminosity.

All of the blue diamonds are well fit by the relation

$$N2 = -0.54 - 0.25(M_B(AB) + 20), \quad (5)$$

which we show as the black solid line in Figure 23. Translating Equation 5 to a metallicity-luminosity relation using the Pettini & Pagel (2004) linear relation gives

$$12 + \log(O/H) = 8.59 - 0.14(M_B(AB) + 20). \quad (6)$$

For $M_B(AB) = -19$ to -20 , the median value of $12 + \log(O/H)$ for the optically-confirmed Ly α Galaxies lies about 0.2 dex lower than that for the combined NUV-continuum selected galaxies. At brighter magnitudes there is less separation, but this is almost certainly an effect of the saturation in $N2$ as the metallicity approaches the solar metallicity (see the right-hand region




FIG. 23.— $[\text{NII}]\lambda 6584/\text{H}\alpha$ ratio vs. absolute rest-frame B magnitude for the *GALEX* sources that appear to be star formers based on their optical and UV spectra and that lie in the redshift interval $z = 0.195 - 0.44$. These include the optically-confirmed $\text{Ly}\alpha$ Galaxies (red solid triangles) and the optically-confirmed NUV-continuum selected galaxies (blue open diamonds). We also show the GOODS-N NUV < 24 galaxy sample with redshifts between $z = 0.15$ and $z = 0.48$ (blue solid diamonds). The error bars are $\pm 1\sigma$. The black line shows the linear fit of $N2 = \log([\text{NII}]\lambda 6584/\text{H}\alpha)$ relative to $M_B(\text{AB})$ for all of the blue diamonds. The metallicity that would be inferred from the Pettini & Pagel (2004) calibration is shown on the right-hand axis. Finally, we show the strong optical emission-line selected galaxies in the redshift interval $z = 0.2 - 0.45$ from the Kakazu et al. (2007) and Hu et al. (2009) USEL samples (black solid squares) and from the Salzer et al. (2009) KISS sample (black open squares).

of Figure 22(a)). A similar effect is seen in the very bright *GALEX* sample of Hoopes et al. (2007).

We also include in Figure 23 strong optical emission-line selected galaxies such as the USELs (black solid squares) of Kakazu et al. (2007) and Hu et al. (2009) and the KISS galaxies (black open squares) of Salzer et al. (2009). The Cardamone et al. (2009) sample of strong emission-line objects (found with a green color selection from the SDSS galaxies) lies closer to the track of the NUV-continuum selected galaxies, and we do not show these on the figure. The optically-confirmed $\text{Ly}\alpha$ Galaxies appear to stretch from the values of the NUV-continuum selected galaxies down to the values of the strong optical emission-line selected galaxies. This appears consistent with the range of $\text{EW}(\text{H}\alpha)$ in the optically-confirmed $\text{Ly}\alpha$ Galaxies (Figure 21), which shows a substantial but not complete overlap with the strong optical emission-line galaxies.

Six of the optically-confirmed $\text{Ly}\alpha$ Galaxies have significantly detected ($> 4\sigma$) $[\text{OIII}]\lambda 4363$ auroral lines. However, one of these (GALEX1240+6233) is classified as an AGN based on the BPT diagram. For the five remaining sources we used the ‘direct’ or T_e method to determine the metallicity (e.g., Seaton 1975; Pagel et al. 1992; Pilyugin & Thuan 2005; Izotov et al. 2006). To derive $T_e[\text{OIII}]$ and the oxygen abundances, we used the Izotov et al. (2006) formulae, which were developed with the latest atomic data and photoionization models. All five sources give abundances which are broadly consistent with the Pettini & Pagel (2004) $N2$ determinations.

The lowest metallicity source is GALEX1417+5228.




FIG. 24.— Measured velocity dispersion σ from the $\text{H}\alpha$ line widths vs. absolute rest-frame B magnitude for the *GALEX* sources that appear to be star formers based on their optical spectra and that lie in the redshift interval $z = 0.195 - 0.44$. These include the optically-confirmed $\text{Ly}\alpha$ Galaxies (red solid triangles) and the optically-confirmed NUV-continuum selected galaxies (blue open diamonds). The error bars are $\pm 1\sigma$. We also show the GOODS-N NUV < 24 galaxy sample with redshifts between $z = 0.15$ and $z = 0.48$ (this excludes broad-line AGNs and intermediate-type Seyferts; blue solid diamonds). Finally, we show the strong optical emission-line selected galaxies in the redshift interval $z = 0.2 - 0.45$ from the Kakazu et al. (2007) and Hu et al. (2009) USEL samples (black solid squares).

This extremely high $\text{EW}(\text{H}\alpha)$ ($\sim 1400 \text{ \AA}$ in the rest frame) source is shown in Figure 5. It is the interesting source we mentioned in Section 5.1. The $[\text{NII}]\lambda 6584/\text{H}\alpha$ ratio in Figure 23 is only at the 1σ level, and the 1σ upper limit from the Pettini & Pagel (2004) calibration gives $12 + \log(\text{O}/\text{H}) < 7.4$. The $[\text{OIII}]\lambda 4363$ line is exceptionally strong, from which we derive $12 + \log(\text{O}/\text{H}) = 7.65 \pm 0.03$. Thus, GALEX1417+5228 is near the extremely low-metallicity class and below all of the strong emission-line sources in the KISS sample of Salzer et al. (2009), as can be seen from Figure 23. The remaining 4 galaxies (GALEX0332-2811, GALEX0959+0151, GALEX1000+0201, and GALEX1001+0233) have direct abundances $12 + \log(\text{O}/\text{H}) = (8.14 \pm 0.03, 8.27 \pm 0.03, 7.96 \pm 0.12 \text{ and } 7.94 \pm 0.03)$ compared to the $N2$ determinations of $12 + \log(\text{O}/\text{H}) = (7.96 \pm 0.02, 8.17 \pm 0.02, 8.20 \pm 0.02 \text{ and } 8.01 \pm 0.03)$.

6.4. Line Widths

There is also a strong correlation between the line widths measured in $\text{H}\alpha$ and luminosity. In Figure 24 we plot the velocity dispersions σ from the $\text{H}\alpha$ line widths versus $M_B(\text{AB})$. The optically-confirmed $\text{Ly}\alpha$ Galaxies (red solid triangles) and the optically-confirmed NUV-continuum selected galaxies (blue open diamonds) show consistent values for both σ and $M_B(\text{AB})$. They are also broadly consistent with the bright end of the GOODS-N NUV < 24 galaxy sample (blue solid diamonds) with any differences being attributable to differences in the photometry. The USELs (black solid squares) lie at the faint end of the distribution. Thus, the optically-confirmed $\text{Ly}\alpha$ Galaxies are being drawn from a population with the same mass to luminosity ratios and kinematical structure as the NUV-continuum selected galaxy population.




FIG. 25.— NUV–Gunn z color vs. NUV magnitude for the *GALEX* sources that appear to be star formers based on their optical spectra and that lie in the redshift interval $z = 0.195 - 0.44$. These include the optically-confirmed Ly α Galaxies (red solid triangles) and the optically-confirmed NUV-continuum selected galaxies (blue open diamonds). We also show the GOODS-N NUV < 24 galaxy sample with redshifts between $z = 0.15$ and $z = 0.48$ (blue solid diamonds).

6.5. Colors and Extinctions

The optically-confirmed Ly α Galaxies also have bluer colors than the optically-confirmed NUV-continuum selected galaxies of the same luminosity. We illustrate this in Figure 25, where we plot observed-frame NUV–Gunn z color using the Gunn z model C magnitudes from the SDSS versus NUV magnitude for the optically-confirmed Ly α Galaxies (red solid triangles) and for the optically-confirmed NUV-continuum selected galaxies (blue open diamonds). While the number of sources in the sample is small, it appears that the Ly α Galaxies are approximately a magnitude bluer in color than the NUV-continuum selected galaxies of the same absolute magnitude. We also show in the figure the GOODS-N NUV < 24 galaxy sample (blue solid diamonds) using the *HST* ACS F850LP AUTO magnitudes. The spread in colors matches the distribution seen in both *GALEX* samples, and we can see that as we move to fainter NUV magnitudes, the fraction of blue sources increases. This may suggest that Ly α Galaxies will be more common in sources below the NUV ~ 22 limit of the *GALEX* spectroscopic observations.

The NUV–Gunn z color difference appears to be at least partly an extinction effect. In Figure 26 we plot the $H\beta/H\alpha$ Balmer ratio versus the NUV–Gunn z color using the Gunn z model C magnitudes from the SDSS. We show only sources where the spectra are of high enough quality to make an accurate measurement and where rest-frame $\text{EW}(H\alpha) < 1000 \text{ \AA}$ to avoid galaxies where the broadband colors are substantially perturbed by the emission lines. However, the emission lines act in the sense of reducing the difference between the populations, since in the stronger $H\alpha$ emission-line galaxies, which are primarily found in the LAE sample, the line contribution brightens the z -band if it is in the correct redshift interval and makes the NUV–Gunn z colors redder. The




FIG. 26.— Balmer ratio $H\beta/H\alpha$ vs. NUV–Gunn z color for the *GALEX* sources that appear to be star formers based on their optical spectra and that lie in the redshift interval $z = 0.195 - 0.44$. These include the optically-confirmed Ly α Galaxies (red solid triangles) and the optically-confirmed NUV-continuum selected galaxies (blue open diamonds). Note that we show only sources where the spectra are of high enough quality to make an accurate measurement and where rest-frame $\text{EW}(H\alpha) < 1000 \text{ \AA}$ to avoid galaxies where the broadband colors are substantially perturbed by the emission lines. The green horizontal line shows the expected case B ratio in the absence of extinction.

Balmer line fluxes are calculated from the observed EWs combined with the continuum fluxes at the emission-line wavelengths inferred from the broadband magnitudes of the galaxies. For $H\beta$ we have applied a $1 \text{ \AA}$ correction to the EWs to allow for underlying stellar absorption (Cowie & Barger 2008). The green horizontal line shows the expected case B ratio in the absence of extinction.

The bluer optically-confirmed Ly α Galaxies are consistent with having only weak extinction within the rather substantial systematic uncertainties, while the redder galaxies, which include the optically-confirmed NUV-continuum selected galaxies, have a lower average Balmer ratio. However, the difference in the Balmer ratios is not statistically significant with a rank sum test giving a 7% probability that the two samples are drawn from the same distribution. The median extinctions inferred from the two samples are $A_v = 1.30$ (0.94, 2.16) for the LAEs and 2.64 (1.61, 2.85) for the optically-confirmed NUV-continuum selected galaxies, where the quantities in brackets give the 68% confidence range. The difference in the color distributions is marginally significant with only a 1.7% probability that the two distributions are the same. However, it is clear that larger samples and preferably more accurate measurements of the Balmer fluxes are required to proceed further.

We may also compare with the UV spectral slopes as measured in the *GALEX* spectra. As can be seen visually in Figure 19, there is very little difference between the average slope of the LAE selected galaxies and the average slope of the NUV-continuum selected galaxies. We have measured the slopes of all of the individual galaxies in the LAE sample (Table 11) and in the NUV-continuum sample (Table 13). The median β of the LAE (NUV-continuum) sample is -1.23 ± 0.22 (-1.07 ± 0.28), and

there appears to be no significant difference in the distribution of the slopes in the two samples. The measured continuum slope translates to an A_{1600} extinction of 1.88 (2.20) for the LAE (NUV-continuum) samples using the calibration of Meurer et al. (1999). (This ignores any small correction for the wavelength range over which the index is measured, which should be similar for the two samples.) Thus, the absolute continuum UV extinction does not appear to be related to the strength of the Ly α line, which must have a complex and indirect dependence on the metallicity and optical colors where there is a significant dependence.

6.6. Galaxy Morphologies

We used the deep i' -band ground-based data from the CFHT MegaPipe database of the GROTH 00 and SIRTFFL 00 fields to make rough morphological classifications of the galaxies in our samples. We divided the galaxies into three classes: spirals, mergers showing clear signs of major interactions, and smaller compact or irregular galaxies. The classifications were made in a blind fashion without reference to the properties of the galaxies to avoid any subjective bias. Most of the galaxies in the optically-confirmed NUV-continuum selected galaxy sample are large, easily recognizable spirals (lower panel of Figure 27). This is also true for the GOODS-N galaxies with $\text{NUV} = 20 - 22$, where all but one of the sources (a merger) fall into this class. The optically-confirmed Ly α Galaxies are much more heterogeneous (upper panel of Figure 27). Perhaps somewhat surprisingly, some of the Ly α Galaxies are large face-on spiral galaxies (see also Finkelstein 2009). However, as can be seen from Figure 28, the Ly α Galaxy sample (red bars) contains a much larger fraction of mergers and compact galaxies than the NUV-continuum selected sample (blue bars).

In Figure 29 we show the BPT diagrams for the optically-confirmed Ly α Galaxy sample (red solid triangles) and for the optically-confirmed NUV-continuum selected sample (blue open diamonds) divided by morphology into (a) spirals, (b) mergers, and (c) compacts. All of the spirals, including the Ly α Galaxies, have high N2 ratios, showing that they are near-solar metallicity sources. The mergers (all three are Ly α Galaxies) appear to have slightly lower metallicities than the spirals. Finally, based on this small sample, the compact galaxies appear to split between higher metallicity NUV-continuum selected galaxies and lower metallicity Ly α Galaxies. Thus, the Ly α Galaxies appear to be a mixture of normal spirals, merging galaxies, and low-metallicity compact and irregular galaxies. It is these latter sources that seem primarily to weight the metallicities of the Ly α Galaxies to lower values than those of the NUV continuum-selected galaxies.

7. CONCLUSIONS

Perhaps the single most significant conclusion we can draw from the low-redshift *GALEX* samples is that LAEs are much less common at low redshifts than they were in the past (Deharvang et al. 2008). In this paper we have shown that formally defined LAEs (rest-frame $\text{EW}(\text{Ly}\alpha) > 20 \text{ \AA}$) constitute about 5% of the local NUV-continuum selected population at $z = 0.3$, as opposed to 20–25% of this population at $z = 3$ (Shapley et al. 2003).




FIG. 27.— The morphologies of some of the *GALEX* sources in the GROTH 00 field that appear to be star formers based on their optical spectra and that lie in the redshift interval $z = 0.195 - 0.44$. These include sources in the optically-confirmed Ly α Galaxy sample (left panel) and sources in the optically-confirmed NUV-continuum selected galaxy sample (right panel). The blue, green, and red colors correspond to the u^* , g' , and i' band images from the CFHT Legacy Survey deep observations of this field. In the right panel nearly all of the galaxies are spirals. We classify only the two right-most galaxies in the left row as compact. Some of the optically-confirmed Ly α Galaxies in the left panel are also spirals (the bottom three galaxies), but a much larger fraction are mergers (the leftmost galaxy in the second row) or compacts (the remaining galaxies).




FIG. 28.— Distribution of galaxy types for the *GALEX* sources in the GROTH 00 and SIRTFFL 00 fields that appear to be star formers based on their optical spectra and that lie in the redshift interval $z = 0.195 - 0.44$. These include the optically-confirmed Ly α Galaxies (red bars) and the optically-confirmed NUV-continuum selected galaxies (blue bars). The NUV-continuum selected galaxies are primarily spirals, while the Ly α Galaxies contain a much larger fraction of mergers and compact galaxies.

This rise is probably best seen by looking at the relative evolution of the volume emissivities of Ly α in LAEs and the volume emissivities of $\nu L_\nu(1500 \text{ \AA})$ in UV-continuum selected galaxies with redshift. The volume emissivities are obtained from the integrated LF fits at each redshift. In Figure 30 we show the Ly α luminosity densities in the LAEs (low redshifts, this paper; higher redshifts, K. Nilsson 2009, priv. comm.; Gronwall et al. 2007; Ouchi et al. 2008) with red solid triangles. We show the UV-continuum luminosity densities from Tresse et al. (2007) with blue solid diamonds. Both rise rapidly from the present time to redshift three, but the rise in the Ly α luminosity densities is much larger than the rise in the UV-continuum luminosity densities. At higher redshifts the UV-continuum luminosity densities turn down while the Ly α luminosity densities remain flat. Thus, as a function of increasing redshift over the whole redshift range $z = 0 - 6$ we appear to be seeing an increase in the amount of escaping Ly α relative to the escaping UV-continuum from the entire UV-continuum selected galaxy population. We note that we are not making an extinction correction in either quantity here and that the comparison is of the light emerging from the galaxy population in the Ly α line and the UV continuum.

The complexity of the low-redshift population makes it hard to provide a single explanation for this evolution. Indeed, the second clear result from the *GALEX* data is that low-redshift Ly α galaxies are not a monolithic population. There is a large fraction of low-metallicity compact galaxies, as might be expected, and merging also seems to make it easier to see Ly α . However, there is also a population of more normal near-solar metallicity spiral galaxies contained in the population. Furthermore, while on average the low-redshift Ly α galaxies have lower metallicities than the NUV-continuum selected galaxies without Ly α emission, the range of metallicities in the




FIG. 29.— BPT diagrams for the *GALEX* sources that appear to be star formers based on their optical spectra, lie in the redshift interval $z = 0.195 - 0.44$, and are in the GROTH 00 or SIRTFFL 00 fields where we have made morphological classifications. Only sources with significant detections ($> 3\sigma$) in either of the H β or [OIII] $\lambda 5007$ lines are shown. The sample is divided into optically-confirmed Ly α Galaxies (red solid triangles) and optically-confirmed NUV-continuum selected galaxies (blue open diamonds) divided by morphology: (a) spirals, (b) mergers, and (c) compacts.




FIG. 30.— Relative evolution of the Ly α luminosity densities in the LAEs (red solid triangles) and the $\nu L_\nu(1500 \text{ \AA})$ luminosity densities in UV-continuum selected galaxies (blue solid diamonds). The Ly α luminosity densities are from the present work at the lowest redshift and from K. Nilsson (2009, priv. comm.), and Ouchi et al. (2008) at higher redshifts. Gronwall et al. (2007) gives a similar luminosity density at $z = 3.1$. The UV-continuum luminosity densities are from Tresse et al. (2007). ± 1 sigma error bars are shown for all the points except that of Nilsson. In some cases they are smaller than the symbol size.




FIG. 31.— $[\text{NII}]\lambda 6584/\text{H}\alpha$ ratio vs. absolute rest-frame B magnitude for the *GALEX* sources that appear to be star formers based on their UV and optical spectra and that lie in the redshift interval $z = 0.195 - 0.44$. These include the optically-confirmed Ly α Galaxies (red solid triangles) and the optically-confirmed NUV-continuum selected galaxies (blue open diamonds). The error bars are $\pm 1\sigma$. We also show the GOODS-N NUV < 24 galaxy sample with redshifts between $z = 0.15$ and $z = 0.48$ (blue solid diamonds). The black line shows the linear fit of $N2 = \log([\text{NII}]\lambda 6584/\text{H}\alpha)$ relative to $M_B(\text{AB})$ for all of the blue diamonds. The metallicity that would be inferred from the Pettini & Pagel (2004) calibration is shown on the right-hand axis. Finally, we show the values measured for the $z \sim 2$ LBGs by Erb et al. (2006; green inverted solid triangles). The green line shows the local metallicity-magnitude relation shifted by 3 magnitudes to match the metallicities of the $z \sim 2$ sources.

Ly α galaxies is wide, stretching from near-solar metallicities down to extremely metal-poor galaxies.

Nevertheless, the overall observed trend of an increasing Ly α fraction with increasing redshift would be ex-

pected if the metals and dust content are decreasing as we move to high redshift. These factors increase the probability of seeing Ly α in the low-redshift sample. The one aspect of this that we can test with the present data is the metallicity evolution. We can make a detailed comparison with high-redshift studies, because Erb et al. (2006) also used the N2 relation in determining the metallicities of their $z \sim 2$ galaxy sample. In Figure 31 we compare our determinations of N2 for the optically-confirmed Ly α Galaxies (red solid triangles), the optically-confirmed NUV-continuum selected galaxies (blue open diamonds), and the GOODS-N NUV-continuum selected galaxies (blue solid diamonds) with the Erb et al. (2006) determinations of N2 for the $z \sim 2$ galaxies (green inverted solid triangles) versus $M_B(\text{AB})$. (Note that the Erb et al. points differ from the low-redshift points in that they are binned averages rather than individual points corresponding to single galaxies.) Consistent with Erb et al. (2006), we find that at the same value of N2, the $z \sim 2$ sources are 3 mag brighter in $M_B(\text{AB})$ (green line) than our $z \sim 0.3$ NUV-continuum selected sample. In translating this into a metallicity, we must remember the dependence of N2 on the ionization parameter. A factor of only 3 increase in q between the typical local value and $z \sim 2$ would increase the inferred metallicity in the $z \sim 2$ galaxies by 0.21 dex and remove much of the inferred evolution. However, in the simplest interpretation, where we treat this purely as a metallicity effect, the $z \sim 2$ galaxies have relatively low metallicities for their luminosities, and they lie in the same metallicity range as the $z \sim 0.3$ Ly α galaxies. We can speculate that a higher fraction of low-metallicity galaxies at $z \sim 2$ might equate to a higher fraction of LAEs at $z \sim 2$, which could be a partial explanation for the higher fraction of LAEs observed at high redshifts than at low redshifts.

We would like to thank the anonymous referee and Jean-Michel Deharvang for critically reading the first draft of this manuscript and for providing extensive and extremely helpful comments. We would also like to thank Kim Nilsson and Steve Finkelstein for their input, which also improved the paper. We are indebted to the staff of the Subaru and Keck Observatories for their excellent assistance with the observations, and to Drew Phillips and Greg Wirth for their help in creating innovative mask designs for the DEIMOS spectrograph. We gratefully acknowledge support from NSF grants AST-0709356 (L. L. C.), AST-0708793 (A. J. B.), and AST-0687850 (E. M. H.), the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation and the David and Lucile Packard Foundation (A. J. B.), and a grant from NASA through an award issued by JPL 1289080 (E. M. H.). Some of the data presented in this paper were obtained from the Multimission Archive at the Space Telescope Science Institute (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-*HST* data is provided by the NASA Office of Space Science via grant NAG5-7584 and by other grants and contracts. This research used the facilities of the Canadian Astronomy Data Centre operated by the National Research Council of Canada with the support of

the Canadian Space Agency. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, Cal-

ifornia Institute of Technology, under contract with the

REFERENCES

- Adelman-McCarthy, J. K., et al. 2008, *ApJS*, 175, 297
- Alexander, D. M., et al. 2003, *AJ*, 126, 539
- Arnouts, S., et al. 2005, *ApJ*, 619, L43
- Atek, H., Kunth, D., Schaerer, D., Hayes, M., Deharvang, J.-M., Östlin, G., & Mas-Hesse, J. M. 2009, *A&A*, 506, L1
- Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, *PASP*, 93, 5 (BPT)
- Barger, A. J., Cowie, L. L., Mushotzky, R. F., Yang, Y., Wang, W.-H., Steffen, A. T., & Capak, P. 2005, *AJ*, 129, 578
- Barger, A. J., Cowie, L. L., & Wang, W.-H. 2008, *ApJ*, 689, 687
- Cardamone, C., et al. 2009, *MNRAS*, 399, 1191
- Cowie, L. L., & Barger, A. J. 2008, *ApJ*, 686, 72
- Cowie, L. L., Barger, A. J., & Trouille, L. 2009, *ApJ*, 692, 1476
- Cowie, L. L., & Hu, E. M. 1998, *AJ*, 115, 1319
- Cowie, L. L., Songaila, A., Hu, E. M., & Cohen, J. G. 1996, *AJ*, 112, 839
- Deharvang, J.-M., et al. 2008, *ApJ*, 680, 1072
- Erb, D. K., Shapley, A. E., Pettini, M., Steidel, C. C., Reddy, N. A., & Adelberger, K. L. 2006, *ApJ*, 644, 813
- Faber, S. M. et al. 2003, *Proc. SPIE*, 4841, 1657
- Felten, J. E. 1976, *ApJ*, 207, 700
- Ferland, G. J., & Osterbrock, D. E. 1985, *ApJ*, 289, 105
- Finkelstein, S. L. 2009, www.iap.fr/col2009/scientific_organisation.html
- Finkelstein, S. L., Cohen, S. H., Malhotra, S., & Rhoads, J. E. 2009, *ApJ*, 700, 276
- Finkelstein, S. L., Rhoads, J. E., Malhotra, S., Pirzkal, N., & Wang, J. 2007, *ApJ*, 660, 1023
- Giavalisco, M., et al. 2004, *ApJ*, 600, L93
- Gronwall, C., et al. 2007, *ApJ*, 667, 79
- Groves, B., Heckman, T., & Kauffmann, G. 2006, *MNRAS*, 371, 1559
- Hoopes, C. G., et al. 2007, *ApJ*, 173, 441
- Hu, E. M., & Cowie, L. L. 2006, *Nature*, 440, 1145
- Hu, E. M., Cowie, L. L., Kakazu, Y., & Barger, A. J. 2009, *ApJ*, 698, 2014
- Hu, E. M., Cowie, L. L., & McMahon, R. G. 1998, *ApJ*, 502, L99
- Izotov, Y. I., Stasińska, G., Meynet, G., Guseva, N. G., & Thuan, T. X. 2006, *A&A*, 448, 955
- Izotov, Y. I., & Thuan, T. X. 2008, *ApJ*, 687, 133
- Kakazu, Y., Hu, E. M., & Cowie, L. L. 2007, *ApJ*, 668, 853
- Kauffmann, G., et al. 2003, *MNRAS*, 346, 1055
- Kennicutt, R. C., Jr. 1998, *ARA&A*, 36, 189
- Kewley, L. J., & Dopita, M. A. 2002, *ApJS*, 142, 35
- Kewley, L. J., Dopita, M. A., Sutherland, R. S., Heisler, C. A., & Trevena, J. 2001, *ApJ*, 556, 121
- Kobulnicky, H. A., & Kewley, L. J. 2004, *ApJ*, 617, 240
- Kunth, D., Leitherer, C., Mas-Hesse, J. M., Östlin, G., & Petrosian, A. 2003, *ApJ*, 597, 263
- Lehmer, B. D., et al. 2005, *ApJS*, 161, 21
- Liang, Y. C., Hammer, F., Flores, H., Gruel, N., & Assemat, F. 2004, *A*, 417, 905
- Leitherer, C. et al. 1999, *ApJS*, 123, 3
- Luo, B., et al. 2008, *ApJS*, 179, 19
- Mallery, R. P. 2009, www.iap.fr/col2009/scientific_organisation.html
- Markwardt, C. B. 2008, *Astronomical Data Analysis Software and Systems XVIII*, Quebec, Canada, ASP Conference Series, Vol. 411, eds. D. Bohlender, P. Dowler and D. Durand Astronomical Society of the Pacific: San Francisco), p. 251-254
- Martin, D. C., et al. 2005, *ApJ*, 619, L1
- McGaugh, S. 1991, *ApJ*, 380, 140
- Meurer, G., Heckman, T. M., and Calzetti, D. 1999, *ApJ*, 521, 64
- More, J. 1978, *Numerical Analysis*, vol. 630, ed. G. A. Watson (Springer-Verlag: Berlin), p. 105
- Morrissey, P., et al. 2007, *ApJS*, 173, 682
- Ouchi, M., et al. 2008, *ApJS*, 176, 301
- Neufeld, D. A. 1991, *ApJ*, 370, L85
- Pagel, B. E. J., Edmunds, M. G., Blackwell, D. E., Chun, M. S., & Smith, G. 1979, *MNRAS*, 189, 95
- Pagel, B. E. J., Simonson, E. A., Terlevich, R. J., & Edmunds, M. G. 1992, *MNRAS*, 255, 325
- Pettini, M., & Pagel, B. E. J. 2004, *MNRAS*, 348, L59
- Pettini, M., Shapley, A. E., Steidel, C. C., Cuby, J.-G., Dickinson, M., Moorwood, A. F. M., Adelberger, K. L., & Giavalisco, M. 2001, *ApJ*, 554, 981
- Pilyugin, L. S., & Thuan, T. X. 2005, *ApJ*, 631, 231
- Prescott, M. K. M., Impey, C. D., Cool, R. J., & Scoville, N. Z. 2006, *ApJ*, 644, 100
- Salpeter, E. E. 1955, *ApJ*, 121, 161
- Salzer, J. J., Williams, A. L., & Gronwall, C. 2009, *ApJ*, 695, L67
- Schaerer, D., & Verhamme, A. 2008, *A&A*, 480, 369
- Schechter, P. 1976, *ApJ*, 203, 297
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Scoville, N., et al. 2007, *ApJS*, 172, 38
- Seaton, M. J. 1975, *MNRAS*, 170, 475
- Shapley, A. E., Steidel, C. C., Pettini, M., & Adelberger, K. L. 2003, *ApJ*, 588, 65
- Siana, B., et al. 2007, *ApJ*, 668, 62
- Storchi-Bergmann, T., Calzetti, D., & Kinney, A. L. 1994, *ApJ*, 429, 572
- Thuan, T. X., & Izotov, Y., 2005, *ApJS*, 161, 240
- Tresse, L., et al. 2007, *A&A*, 472, 403
- Tresse, L., & Maddox, S. J. 1998, *ApJ*, 495, 691
- Trouille, L., Barger, A. J., Cowie, L. L., Yang, Y., & Mushotzky, R. F. 2008, *ApJS*, 179, 1
- Trouille, L., Barger, A. J., Cowie, L. L., Yang, Y., & Mushotzky, R. F. 2009, *ApJ*, 703, 2160
- van Breukelen, C., Jarvis, M. J., & Venemans, B. P. 2005, *MNRAS*, 359, 895
- Virani, S. N., Treister, E., Urry, C. M., & Gawiser, E. 2006, *AJ*, 131, 2373
- Yencho, B., Barger, A. J., Trouille, L., & Winter, L. M. 2009, *ApJ*,

TABLE 1
GALEX FIELDS

Name (1)	R.A. (J2000.0) (2)	Decl. (J2000.0) (3)	Exposure (4)	NUV _{lim} (5)	E(B-V) (6)	Gal Lat (7)	Number (8)	Number(ID) (9)
GROTH 00	214.99182	52.78173	291	21.8	0.007	59.5	1152	69
NGPDWS 00	219.15610	35.17135	156	21.5	0.009	66.2	738	46
CDFS 00	53.12779	-27.87137	149	21.5	0.008	-54.4	876	39
COSMOS 00	150.11900	2.20583	140	21.5	0.018	42.1	779	29
ELAISS1 00	9.63857	-43.99023	84	21.2	0.007	-72.9	589	19
SIRTF 00	259.12387	59.90915	80	21.2	0.020	35.0	800	22
LOCK 00	162.67843	58.73117	48	20.9	0.009	52.1	390	12
SIRTF 01	260.41425	59.34286	34	20.7	0.029	34.4	612	14
HDFN 00	189.20946	62.19772	24	20.5	0.012	54.8	265	10

TABLE 2
GALEX SPECTRAL SAMPLE IDENTIFICATIONS: GROTH 00

Number (1)	Name (2)	R.A. (J2000.0) (3)	Decl. (J2000.0) (4)	NUV (5)	FUV (6)	z (7)	Line Width (8)	ID (9)
1	GALEX1416+5252	214.147766	52.871555	21.12	24.14	1.146	9391± 466	AGN
2	GALEX1416+5259	214.201279	52.984028	20.68	21.45	1.233	6221± 177	AGN
3	GALEX1416+5250	214.208328	52.838863	21.49	21.97	0.288	1832± 404	Galaxy
4	GALEX1417+5305	214.291290	53.086723	21.29	21.82	0.268	1985± 227	Galaxy
5	GALEX1417+5235	214.300888	52.599110	21.24	21.76	0.262	1932± 285	AGN
6	GALEX1417+5230	214.352707	52.507141	20.37	21.20	0.896	7618± 554	Galaxy
7	GALEX1417+5230	214.399918	52.508358	20.82	22.39	0.988	6585± 94	AGN
8	GALEX1417+5304	214.419083	53.079052	21.56	22.14	0.271	1587± 281	Galaxy
9	GALEX1417+5232	214.428452	52.538414	21.46	21.84	0.199	3718± 817	Galaxy
10	GALEX1417+5228	214.430923	52.468307	21.90	21.77	0.210	2051± 84	Galaxy
11	GALEX1417+5246	214.438126	52.771694	21.83	22.08	0.248	2719± 326	Galaxy
12	GALEX1417+5303	214.446213	53.063831	21.59	22.53	0.916	4845± 171	AGN
13	GALEX1417+5242	214.474335	52.700500	21.71	22.04	0.205	1890± 531	Galaxy
14	GALEX1418+5252	214.506790	52.866917	20.58	21.71	1.160	9640± 304	AGN
15	GALEX1418+5237	214.518997	52.629444	21.32	22.95	1.090	5943± 214	AGN
16	GALEX1418+5245	214.521545	52.752110	21.29	21.63	0.245	2464± 260	Galaxy
17	GALEX1418+5301(o)	214.563675	53.027248	21.33	24.41	0.988	4657± 210	Galaxy
18	GALEX1418+5217	214.588928	52.287388	21.11	21.60	0.360	4320± 306	AGN
19	GALEX1418+5257(o)	214.589371	52.957279	20.02	20.86	0.389	5093± 243	AGN
20	GALEX1418+5218	214.593506	52.306747	21.70	22.28	0.239	1430± 187	Galaxy
21	GALEX1418+5222	214.626038	52.370251	21.32	22.49	0.824	6734± 219	AGN
22	GALEX1418+5310	214.658005	53.171585	21.01	21.89	0.896	6260± 212	AGN
23	GALEX1418+5224	214.660172	52.400002	19.09	22.41	1.120	10257± 108	AGN
24	GALEX1418+5217	214.701202	52.299168	21.65	21.97	0.242	2586± 448	Galaxy
25	GALEX1418+5307	214.728928	53.130028	22.09	21.89	0.205	2015± 145	Galaxy
26	GALEX1418+5249	214.732056	52.824558	21.11	21.08	0.268	1889± 317	Galaxy
27	GALEX1418+5259	214.733200	52.992195	21.38	21.88	0.288	2652± 287	Galaxy
28	GALEX1418+5218	214.748795	52.302834	20.96	23.86	0.883	5613± 201	AGN
29	GALEX1419+5221	214.780151	52.352386	21.39	22.15	0.268	3167± 462	Galaxy
30	GALEX1419+5315	214.794739	53.265835	20.98	21.90	0.265	2835± 251	Galaxy
31	GALEX1419+5223	214.811249	52.390808	21.77	22.16	0.257	2321± 446	Galaxy
32	GALEX1419+5230	214.907211	52.507057	21.18	21.75	0.285	2314± 400	Galaxy
33	GALEX1419+5213	214.940002	52.228359	21.10	21.99	0.697	10281±1660	AGN
34	GALEX1419+5221	214.970749	52.350250	21.29	21.83	0.268	2906± 456	Galaxy
35	GALEX1420+5216	215.034515	52.279945	18.73	20.26	1.060	10064± 93	AGN
36	GALEX1420+5316	215.099625	53.268028	20.54	21.91	0.740	9933± 380	AGN
37	GALEX1420+5247(b)	215.132080	52.799389	20.73	21.10	0.257	2388± 163	Galaxy
38	GALEX1420+5308(o)	215.154327	53.144749	21.12	21.93	0.202	4019± 825	Galaxy
39	GALEX1420+5306(b)	215.170456	53.114002	21.05	21.46	0.199	1883± 357	Galaxy
40	GALEX1420+5217	215.174042	52.283916	21.42	22.99	1.017	6146± 571	Galaxy
41	GALEX1420+5243	215.180359	52.718750	21.29	21.54	0.251	2756± 291	Galaxy
42	GALEX1420+5236	215.181488	52.603306	20.89	21.36	0.340	4820± 195	AGN
43	GALEX1420+5250	215.186218	52.835056	21.36	21.63	0.254	2381± 262	Galaxy
44	GALEX1420+5256	215.218536	52.939640	18.28	19.00	0.683	10794± 111	AGN
45	GALEX1420+5248	215.236969	52.808361	20.89	22.00	1.201	8131± 355	AGN
46	GALEX1420+5313	215.237045	53.218750	21.26	23.87	0.749	3732± 371	Galaxy
47	GALEX1421+5241	215.301178	52.696415	20.82	21.58	0.844	8319± 294	AGN
48	GALEX1421+5238	215.341003	52.634724	21.27	21.91	0.282	3897± 296	Galaxy
49	GALEX1421+5239	215.352737	52.655472	21.82	22.23	0.257	2055± 264	Galaxy
50	GALEX1421+5305	215.372772	53.097027	21.53	23.83	1.190	7712± 552	AGN
51	GALEX1421+5302	215.382706	53.044975	21.53	22.58	0.965	5377± 299	AGN
52	GALEX1421+5308	215.387955	53.136333	21.82	22.26	0.202	1252± 0	Galaxy
53	GALEX1421+5234	215.392624	52.574860	21.48	24.06	1.201	7096± 336	AGN
54	GALEX1421+5231	215.399704	52.527527	20.06	20.31	0.251	4066± 93	AGN
55	GALEX1421+5221	215.421539	52.359886	21.64	999.00	0.947	3813± 221	Galaxy
56	GALEX1421+5249	215.463165	52.830891	21.25	21.87	0.202	3119± 445	Galaxy
57	GALEX1421+5224	215.477295	52.406723	20.39	21.22	0.352	2547± 246	Galaxy
58	GALEX1422+5310	215.500092	53.168472	21.65	22.26	0.239	1578± 438	Galaxy
59	GALEX1422+5252	215.533752	52.873722	21.30	22.52	0.305	2056± 130	Galaxy
60	GALEX1422+5305	215.538086	53.099976	19.95	21.16	0.760	8935± 146	AGN
61	GALEX1422+5302	215.571579	53.036446	20.82	22.45	0.904	7143± 171	AGN
62	GALEX1422+5230	215.610550	52.507526	21.56	21.93	0.222	2149± 272	AGN
63	GALEX1423+5233	215.757919	52.554806	20.55	21.58	0.847	7568± 284	AGN
64	GALEX1423+5246	215.767471	52.774918	18.85	20.50	0.999	8139± 59	AGN
65	GALEX1423+5246	215.775833	52.779697	21.33	22.01	0.346	2199± 315	Galaxy
66	GALEX1423+5244	215.787262	52.743389	999.00	20.72	0.280	3365± 33	AGN
67	GALEX1423+5242	215.823959	52.713528	21.02	21.76	0.280	3883± 534	Galaxy
68	GALEX1423+5242	215.835556	52.710999	21.65	24.49	0.757	6831± 864	Galaxy
69	GALEX1423+5244	215.842850	52.742474	21.74	22.35	0.280	2624± 569	Galaxy

TABLE 3
GALEX SPECTRAL SAMPLE IDENTIFICATIONS: NGPDWS 00

Number (1)	Name (2)	R.A. (J2000.0) (3)	Decl. (J2000.0) (4)	NUV (5)	FUV (6)	z (7)	Line Width (8)	ID (9)
1	GALEX1434+3512	218.502533	35.205639	21.15	22.97	1.193	8546± 511	AGN
2	GALEX1434+3502	218.602798	35.044998	20.10	20.19	0.280	4605± 109	AGN
3	GALEX1434+3448	218.672165	34.808582	20.79	21.68	0.732	8811± 432	AGN
4	GALEX1434+3529	218.686600	35.488083	20.08	20.98	0.873	9495± 283	AGN
5	GALEX1434+3456	218.698334	34.934277	19.82	22.97	0.927	9631± 195	AGN
6	GALEX1434+3524	218.710663	35.409168	20.24	21.51	0.994	5629± 85	AGN
7	GALEX1434+3532	218.718338	35.547527	21.44	21.81	0.198	1781± 340	Galaxy
8	GALEX1434+3508	218.739197	35.141582	20.46	22.35	1.213	4443± 316	Galaxy
9	GALEX1435+3524	218.753174	35.408611	21.38	21.95	0.995	5094± 478	Galaxy
10	GALEX1435+3449	218.764587	34.821220	21.32	22.74	0.378	2016± 577	Galaxy
11	GALEX1435+3457	218.781403	34.956833	20.65	22.10	1.117	7680± 171	AGN
12	GALEX1435+3504	218.868118	35.075916	20.32	21.55	1.147	11538± 448	AGN
13	GALEX1435+3445	218.996429	34.751472	21.13	999.00	1.133	5911± 158	AGN
14	GALEX1436+3527	219.026245	35.458553	21.38	21.57	0.252	2477± 254	Galaxy
15	GALEX1436+3532	219.064285	35.549942	21.24	22.21	0.894	6720± 330	AGN
16	GALEX1436+3537	219.074417	35.624084	19.62	22.72	1.078	5673± 215	AGN
17	GALEX1436+3440	219.086914	34.672001	21.09	22.01	0.374	1948± 525	Galaxy
18	GALEX1436+3456	219.092041	34.942139	21.16	21.47	0.272	1716± 141	Galaxy
19	GALEX1436+3459	219.100540	34.993530	21.31	21.57	0.215	2050± 195	Galaxy
20	GALEX1436+3537	219.101288	35.619305	18.87	19.73	0.768	8344± 83	AGN
21	GALEX1436+3525	219.102737	35.427055	19.21	20.06	1.065	11452± 165	AGN
22	GALEX1436+3453	219.131912	34.890221	20.33	21.69	1.068	7087± 132	AGN
23	GALEX1436+3438	219.133331	34.641445	21.39	22.38	0.357	3319± 642	Galaxy
24	GALEX1436+3450	219.152740	34.842945	20.66	21.15	0.283	3280± 184	AGN
25	GALEX1436+3520	219.156403	35.337666	21.26	22.27	1.162	4573± 598	Galaxy
26	GALEX1436+3525	219.173416	35.426388	21.02	21.87	1.028	10346± 957	Galaxy
27	GALEX1436+3539	219.175705	35.658165	19.22	20.41	0.845	11197± 176	AGN
28	GALEX1436+3525	219.193008	35.417641	21.22	22.19	0.370	2450± 321	Galaxy
29	GALEX1436+3439	219.197495	34.650475	21.05	22.34	1.243	6190± 328	AGN
30	GALEX1436+3535	219.243362	35.597694	20.69	21.46	0.208	1481± 248	Galaxy
31	GALEX1437+3508	219.255585	35.137722	20.74	21.41	0.880	8070± 418	AGN
32	GALEX1437+3541	219.281967	35.692280	20.92	22.76	0.664	3050± 374	Galaxy
33	GALEX1437+3534(o)	219.324371	35.580055	21.34	21.77	0.429	4603± 566	AGN
34	GALEX1437+3445	219.334000	34.757637	21.18	21.51	0.328	2007± 147	Galaxy
35	GALEX1437+3448	219.345459	34.801918	20.98	22.04	0.950	8644± 361	AGN
36	GALEX1437+3522	219.347076	35.374138	21.15	999.00	1.061	10026± 608	AGN
37	GALEX1437+3441	219.356308	34.685585	21.00	21.58	0.294	2388± 314	Galaxy
38	GALEX1437+3522	219.388214	35.370251	20.64	21.08	0.247	1694± 286	Galaxy
39	GALEX1437+3500(o)	219.406250	35.011749	21.43	23.15	0.385	1449± 346	Galaxy
40	GALEX1437+3442	219.431671	34.715027	20.02	21.00	1.115	8691± 188	AGN
41	GALEX1437+3528	219.437714	35.473278	21.38	21.91	0.398	2626± 217	AGN
42	GALEX1438+3458	219.523636	34.980556	19.28	19.54	0.431	5134± 194	AGN
43	GALEX1438+3446	219.541245	34.768166	20.00	21.06	0.833	9702± 246	AGN
44	GALEX1438+3518	219.603546	35.305168	20.93	21.63	0.874	6915± 683	Galaxy
45	GALEX1438+3513	219.704224	35.232277	21.14	22.18	0.805	7144± 368	AGN
46	GALEX1438+3457	219.746078	34.960636	21.29	21.88	0.374	1897± 332	Galaxy

TABLE 4
GALEX SPECTRAL SAMPLE IDENTIFICATIONS: CDFS 00

Number (1)	Name (2)	R.A. (J2000.0) (3)	Decl. (J2000.0) (4)	NUV (5)	FUV (6)	z (7)	Line Width (8)	ID (9)
1	GALEX0330-2748	52.624874	-27.815527	21.21	23.94	0.855	5966 $\pm$ 454	AGN
2	GALEX0330-2744	52.666042	-27.747833	21.32	21.96	0.710	9635 $\pm$ 1603	AGN
3	GALEX0330-2735	52.676041	-27.585638	21.10	22.43	1.130	9888 $\pm$ 609	AGN
4	GALEX0330-2759	52.696835	-27.988083	21.15	22.23	0.685	8226 $\pm$ 959	AGN
5	GALEX0330-2732	52.732708	-27.547722	20.41	21.18	1.013	5587 $\pm$ 328	Galaxy
6	GALEX0330-2816	52.737583	-28.279501	21.11	21.55	0.285	2124 $\pm$ 286	Galaxy
7	GALEX0331-2751	52.759998	-27.858416	21.50	21.97	0.340	2769 $\pm$ 375	Galaxy
8	GALEX0331-2756	52.836498	-27.946917	21.30	22.24	0.685	8353 $\pm$ 748	AGN
9	GALEX0331-2816	52.849792	-28.272055	20.92	21.02	0.219	4499 $\pm$ 250	AGN
10	GALEX0331-2811	52.962204	-28.189028	20.68	21.02	0.216	1739 $\pm$ 163	Galaxy
11	GALEX0331-2814	52.976460	-28.238554	21.49	21.84	0.282	2108 $\pm$ 125	Galaxy
12	GALEX0331-2809	52.999332	-28.164444	21.18	21.43	0.239	1415 $\pm$ 174	Galaxy
13	GALEX0332-2801	53.049873	-28.025000	21.51	21.71	0.219	1972 $\pm$ 138	Galaxy
14	GALEX0332-2811	53.061497	-28.186584	21.18	21.71	0.265	2323 $\pm$ 300	Galaxy
15	GALEX0332-2813	53.077999	-28.222332	21.17	21.42	0.282	3232 $\pm$ 172	Galaxy
16	GALEX0332-2740	53.110252	-27.676527	20.04	21.14	1.040	8199 $\pm$ 204	AGN
17	GALEX0332-2741	53.112419	-27.684639	19.82	21.18	0.740	8440 $\pm$ 186	AGN
18	GALEX0332-2808(o)	53.155373	-28.146307	-999.00	24.36	0.775	16316 $\pm$ 0	AGN
19	GALEX0332-2739	53.158833	-27.662472	20.88	22.03	0.838	6348 $\pm$ 207	AGN
20	GALEX0332-2811	53.174255	-28.190195	20.27	20.17	0.208	2070 $\pm$ 78	Galaxy
21	GALEX0332-2810	53.184128	-28.174416	21.04	21.83	1.152	7915 $\pm$ 379	AGN
22	GALEX0332-2822	53.191792	-28.375555	20.32	21.94	0.844	5287 $\pm$ 79	Galaxy
23	GALEX0332-2753	53.236042	-27.887835	20.62	21.10	0.375	1873 $\pm$ 362	Galaxy
24	GALEX0332-2823	53.240330	-28.388361	20.83	21.30	0.219	1739 $\pm$ 154	Galaxy
25	GALEX0333-2821	53.258461	-28.357695	21.08	21.26	0.251	1792 $\pm$ 228	Galaxy
26	GALEX0333-2813	53.268082	-28.227194	20.88	22.75	1.015	5020 $\pm$ 617	Galaxy
27	GALEX0333-2744	53.280293	-27.742332	20.27	20.79	0.220	1567 $\pm$ 293	Galaxy
28	GALEX0333-2725	53.352165	-27.430637	19.64	21.21	1.140	8379 $\pm$ 123	AGN
29	GALEX0333-2727	53.359249	-27.454306	21.17	21.61	0.352	3514 $\pm$ 409	Galaxy
30	GALEX0333-2756	53.370499	-27.944778	21.02	22.32	0.840	7906 $\pm$ 326	AGN
31	GALEX0333-2822	53.386833	-28.372278	20.33	21.65	0.861	6758 $\pm$ 250	Galaxy
32	GALEX0333-2817	53.414001	-28.289944	20.10	21.19	0.991	6436 $\pm$ 140	AGN
33	GALEX0334-2756	53.517128	-27.941610	21.02	22.12	0.988	9311 $\pm$ 411	AGN
34	GALEX0334-2807	53.530209	-28.119057	20.59	22.09	1.080	9411 $\pm$ 559	AGN
35	GALEX0334-2743	53.534500	-27.727251	20.38	21.36	1.028	7839 $\pm$ 197	AGN
36	GALEX0334-2815	53.541370	-28.255472	21.57	22.77	0.345	2339 $\pm$ 444	Galaxy
37	GALEX0334-2745	53.570415	-27.751307	20.19	23.38	1.163	6882 $\pm$ 153	AGN
38	GALEX0334-2737	53.602249	-27.632418	20.43	20.92	0.372	4305 $\pm$ 874	Galaxy
39	GALEX0334-2759	53.662498	-27.987610	21.40	22.01	0.870	6052 $\pm$ 538	Galaxy

TABLE 5
GALEX SPECTRAL SAMPLE IDENTIFICATIONS: COSMOS 00

Number (1)	Name (2)	R.A. (J2000.0) (3)	Decl. (J2000.0) (4)	NUV (5)	FUV (6)	z (7)	Line Width (8)	ID (9)
1	GALEX0958+0205	149.663834	2.085083	21.22	21.72	1.218	5888± 396	AGN
2	GALEX0958+0213	149.739044	2.220722	20.78	23.70	1.025	6963± 191	AGN
3	GALEX0959+0219	149.761612	2.318472	19.99	20.55	0.349	4751± 137	AGN
4	GALEX0959+0209	149.828262	2.164305	21.00	999.00	1.158	4174± 175	AGN
5	GALEX0959+0149	149.873871	1.832972	21.42	21.76	0.208	1378± 320	Galaxy
6	GALEX0959+0212	149.912643	2.200389	21.20	22.07	0.698	5802± 843	AGN
7	GALEX0959+0151	149.917953	1.856028	21.10	21.47	0.254	2662± 313	Galaxy
8	GALEX1000+0235	150.053833	2.589667	19.39	20.66	0.706	9088± 197	AGN
9	GALEX1000+0158	150.105347	1.981167	18.97	19.34	0.379	4161± 58	AGN
10	GALEX1000+0157	150.115784	1.951139	20.72	20.69	0.268	2108± 65	Galaxy
11	GALEX1000+0152	150.139252	1.876972	21.08	23.96	0.832	13437±1614	AGN
12	GALEX1000+0202	150.145676	2.043139	21.25	22.45	1.178	5348± 296	AGN
13	GALEX1000+0201	150.149094	2.020583	20.71	21.03	0.271	2146± 162	Galaxy
14	GALEX1000+0206	150.179825	2.110333	20.46	21.95	0.372	5340± 547	AGN
15	GALEX1000+0152	150.208389	1.875472	21.15	22.44	1.150	9290± 527	AGN
16	GALEX1001+0240(b)	150.262878	2.671028	999.00	21.47	0.268	4056± 282	Galaxy
17	GALEX1001+0220	150.294846	2.346972	21.47	21.51	0.251	1743± 82	Galaxy
18	GALEX1001+0236	150.318039	2.602167	21.36	22.54	0.958	7987± 485	AGN
19	GALEX1001+0227(b)	150.326996	2.460861	21.42	22.23	1.042	3238± 289	Galaxy
20	GALEX1001+0145	150.396637	1.756500	21.50	21.74	0.219	1565± 293	Galaxy
21	GALEX1001+0233	150.413330	2.563611	21.00	21.43	0.389	2361± 367	Galaxy
22	GALEX1001+0202	150.443756	2.049222	20.37	20.99	0.677	5179± 203	AGN
23	GALEX1001+0214	150.449707	2.246444	20.63	21.43	0.881	5863± 256	AGN
24	GALEX1001+0200	150.463043	2.009111	20.33	21.30	0.971	8044± 206	AGN
25	GALEX1002+0224(o)	150.511993	2.409750	20.41	23.54	0.999	10382± 946	Galaxy
26	GALEX1002+0230	150.544846	2.507444	19.87	22.30	1.169	5494± 81	AGN
27	GALEX1002+0229	150.572754	2.499972	21.16	22.78	1.109	6344± 285	AGN
28	GALEX1002+0207	150.579086	2.132917	21.36	22.07	0.375	3140± 381	Galaxy
29	GALEX1002+0156(b)	150.579407	1.933667	20.48	21.11	0.305	3577± 461	Galaxy

TABLE 6
GALEX SPECTRAL SAMPLE IDENTIFICATIONS: ELAISS1 00

Number (1)	Name (2)	R.A. (J2000.0) (3)	Decl. (J2000.0) (4)	NUV (5)	FUV (6)	z (7)	Line Width (8)	ID (9)
1	GALEX0036-4348	9.061083	-43.816639	21.15	22.22	0.876	8611± 598	AGN
2	GALEX0037-4347	9.265166	-43.799919	20.68	23.65	1.240	6359± 362	Galaxy
3	GALEX0037-4342	9.337708	-43.711136	20.61	20.98	0.225	2130± 97	AGN
4	GALEX0037-4338	9.371167	-43.635670	20.52	20.85	0.300	2846± 380	Galaxy
5	GALEX0038-4414	9.511625	-44.233448	20.99	22.21	0.878	5255± 302	AGN
6	GALEX0038-4352	9.520541	-43.874584	20.63	21.08	0.222	2348± 196	Galaxy
7	GALEX0038-4352	9.539542	-43.878113	19.49	20.23	0.936	10019± 258	AGN
8	GALEX0038-4414	9.559041	-44.243584	21.29	21.14	0.271	2211± 79	AGN
9	GALEX0038-4343	9.666250	-43.722443	20.94	21.15	0.210	1846± 334	Galaxy
10	GALEX0039-4341	9.849500	-43.699444	21.13	22.39	0.996	7745± 316	AGN
11	GALEX0039-4417	9.930417	-44.295277	21.18	25.48	1.158	4560± 308	AGN
12	GALEX0039-4341	9.958793	-43.694973	21.10	21.25	0.409	4684± 450	AGN
13	GALEX0039-4405	9.975541	-44.086029	21.06	21.46	0.334	3258± 162	AGN
14	GALEX0040-4355	10.107124	-43.927280	21.00	21.32	0.671	2797± 620	Galaxy
15	GALEX0040-4340	10.117126	-43.671642	19.81	20.43	0.873	8651± 253	AGN
16	GALEX0040-4338	10.172042	-43.638496	20.32	21.06	0.760	6983± 238	AGN
17	GALEX0040-4403	10.177958	-44.061306	19.58	22.14	0.855	8977± 143	AGN
18	GALEX0040-4409	10.243834	-44.164085	20.51	21.16	1.233	5461± 301	AGN
19	GALEX0041-4352	10.272666	-43.874641	20.64	21.54	0.311	3413± 573	Galaxy

TABLE 7
GALEX SPECTRAL SAMPLE IDENTIFICATIONS: SIRTFFL 00

Number (1)	Name (2)	R.A. (J2000.0) (3)	Decl. (J2000.0) (4)	NUV (5)	FUV (6)	z (7)	Line Width (8)	ID (9)
1	GALEX1712+6007	258.131622	60.122082	20.98	999.00	1.204	5908± 388	AGN
2	GALEX1712+6005	258.144928	60.091888	19.55	20.85	0.999	9189± 215	AGN
3	GALEX1712+5956	258.149719	59.946888	20.37	20.99	0.225	1296± 243	Galaxy
4	GALEX1712+6001	258.197723	60.022778	20.80	23.63	0.786	3812± 316	Galaxy
5	GALEX1713+6001	258.290100	60.024193	20.26	21.42	0.818	6461± 239	AGN
6	GALEX1714+5949	258.591766	59.833332	21.09	21.28	0.231	1946± 179	Galaxy
7	GALEX1714+5956	258.683746	59.947247	21.13	21.62	0.222	2059± 482	Galaxy
8	GALEX1715+5956	258.751678	59.936642	19.71	21.07	0.695	12588±1511	AGN
9	GALEX1715+5946	258.807953	59.777473	20.05	20.22	0.251	3441± 108	AGN
10	GALEX1716+5932	259.178070	59.548889	21.04	21.79	0.360	1574± 501	Galaxy
11	GALEX1717+5925	259.358429	59.433167	20.92	21.84	0.953	8569± 598	AGN
12	GALEX1717+5929	259.379578	59.487473	20.34	20.94	0.300	1818± 318	Galaxy
13	GALEX1717+5947	259.387543	59.796555	19.40	20.58	1.060	11329± 331	AGN
14	GALEX1717+5944	259.436890	59.744305	20.79	22.12	0.939	10577± 760	Galaxy
15	GALEX1717+5932	259.448639	59.549419	19.28	19.42	0.251	9213± 241	AGN
16	GALEX1717+5944	259.465027	59.748554	21.08	21.66	0.196	1335± 395	Galaxy
17	GALEX1718+5933	259.528839	59.553638	20.15	20.34	0.274	8108± 279	AGN
18	GALEX1718+5933	259.666138	59.566586	18.49	18.86	0.381	9816± 256	AGN
19	GALEX1718+6023	259.671295	60.399696	20.22	20.95	0.657	10217±1980	AGN
20	GALEX1718+5949	259.748260	59.825752	20.91	23.31	0.372	1401± 291	Galaxy
21	GALEX1719+5950	259.752808	59.845055	20.58	21.93	0.971	13300±1287	Galaxy
22	GALEX1719+6013	259.963531	60.222721	19.65	21.33	1.060	10279± 273	AGN

TABLE 8
GALEX SPECTRAL SAMPLE IDENTIFICATIONS: LOCK 00

Number (1)	Name (2)	R.A. (J2000.0) (3)	Decl. (J2000.0) (4)	NUV (5)	FUV (6)	z (7)	Line Width (8)	ID (9)
1	GALEX1047+5827	161.773422	58.461693	20.33	21.13	0.248	4451± 478	Galaxy
2	GALEX1048+5910	162.048798	59.179695	20.60	21.46	0.711	7525± 664	AGN
3	GALEX1049+5853	162.371796	58.893944	20.30	20.74	0.251	3975± 205	AGN
4	GALEX1050+5819	162.500885	58.317833	18.04	18.88	0.835	8654± 91	AGN
5	GALEX1051+5838	162.771729	58.642475	20.65	21.35	0.915	4969± 782	AGN
6	GALEX1051+5916	162.775833	59.273808	19.13	20.70	0.775	9575± 222	AGN
7	GALEX1051+5832	162.807800	58.544777	20.89	22.03	0.863	8187± 725	AGN
8	GALEX1051+5913	162.921417	59.218193	19.06	19.72	0.440	7587± 359	AGN
9	GALEX1052+5851	163.072922	58.856194	20.57	21.62	0.970	9599± 628	AGN
10	GALEX1053+5905	163.284698	59.089584	19.15	19.53	0.434	3642± 187	AGN
11	GALEX1053+5855	163.292374	58.925808	19.11	19.98	1.180	6607± 124	AGN
12	GALEX1054+5852	163.587448	58.871887	20.42	21.62	1.184	6321± 353	AGN

TABLE 9
GALEX SPECTRAL SAMPLE IDENTIFICATIONS: SIRTFFL 01

Number (1)	Name (2)	R.A. (J2000.0) (3)	Decl. (J2000.0) (4)	NUV (5)	FUV (6)	z (7)	Line Width (8)	ID (9)
1	GALEX1718+5933	259.665924	59.566555	18.49	18.95	0.396	8589± 398	AGN
2	GALEX1719+5934	259.819031	59.580418	20.04	20.61	0.363	3107± 862	Galaxy
3	GALEX1720+5932	260.134705	59.535336	19.96	20.63	1.169	12141± 862	AGN
4	GALEX1720+5910	260.234222	59.166832	20.46	21.98	1.022	5963± 525	Galaxy
5	GALEX1721+5924	260.269806	59.414192	19.71	20.52	0.783	9317± 490	AGN
6	GALEX1722+5901	260.734222	59.027248	20.56	21.73	0.933	7905± 559	Galaxy
7	GALEX1723+5951	260.793243	59.851692	18.86	21.48	0.996	14026± 680	AGN
8	GALEX1723+5853	260.794189	58.884003	20.63	22.04	1.045	9044± 562	AGN
9	GALEX1723+5936	260.819458	59.601192	20.22	20.35	0.421	2971± 242	AGN
10	GALEX1723+5944	260.847595	59.748249	20.36	20.91	0.375	1346± 811	Galaxy
11	GALEX1724+5936	261.058838	59.612278	20.35	22.28	0.752	8007± 503	AGN
12	GALEX1724+5917	261.112488	59.289333	20.06	20.42	0.257	2371± 216	AGN
13	GALEX1724+5921	261.214172	59.359165	20.40	22.06	1.077	11255± 906	AGN
14	GALEX1725+5920	261.344940	59.345528	20.01	20.35	0.386	1180± 384	Galaxy

TABLE 10
GALEX SPECTRAL SAMPLE IDENTIFICATIONS: HDFN 00

Number (1)	Name (2)	R.A. (J2000.0) (3)	Decl. (J2000.0) (4)	NUV (5)	FUV (6)	z (7)	Line Width (8)	ID (9)
1	GALEX1234+6233	188.504761	62.554085	19.91	20.73	1.123	12033± 655	AGN
2	GALEX1234+6226	188.561661	62.444443	17.99	18.89	0.973	4311± 212	AGN
3	GALEX1234+6220	188.670288	62.335999	20.37	20.85	0.398	2708± 374	AGN
4	GALEX1237+6217	189.279007	62.284000	20.09	20.88	1.022	11390± 878	AGN
5	GALEX1237+6211	189.498337	62.184139	19.41	20.70	0.913	8273± 227	AGN
6	GALEX1238+6202	189.567291	62.035610	19.30	20.21	1.001	10252± 432	AGN
7	GALEX1238+6143	189.702667	61.731861	18.34	19.25	0.832	5570± 66	AGN
8	GALEX1239+6206	189.881638	62.105251	19.75	20.70	1.201	7959± 410	AGN
9	GALEX1240+6233	190.041428	62.561668	20.28	20.69	0.210	2094± 209	Galaxy
10	GALEX1240+6223	190.138916	62.390915	20.44	21.43	1.112	8787±1011	AGN

TABLE 11
DEIMOS OBSERVATIONS: GALEX CANDIDATE Ly α GALAXIES

NAME (1)	R.A. (J2000.0) (2)	Decl. (J2000.0) (3)	NUV (4)	g (5)	z_{galax} (6)	$z_{optical}$ (7)	EW(H α) (8)	Optical Type (9)
GALEX0038-4352	9.520541	-43.874580	20.63	...	0.2220	0.2191	15± 1	Galaxy
GALEX0038-4343	9.666203	-43.722527	20.94	...	0.2104	0.2072	314± 2	Galaxy
GALEX0331-2811	52.962204	-28.189028	20.68	...	0.2162	0.2132	3± 0	Galaxy
GALEX0331-2814	52.976501	-28.238640	21.49	...	0.2824	0.2802	38± 0	Galaxy
GALEX0331-2809	52.999294	-28.164413	21.18	...	0.2392	0.2362	227± 1	Galaxy
GALEX0332-2801	53.049751	-28.025028	21.51	...	0.2191	0.2155	472± 1	Galaxy
GALEX0332-2811	53.061584	-28.186525	21.18	...	0.2651	0.2788	234± 1	Galaxy
GALEX0332-2813	53.078033	-28.222441	21.17	...	0.2824	0.2788	... 1	Seyfert 1.8
GALEX0332-2811	53.174255	-28.190306	20.27	...	0.2076	0.2044	406± 1	Galaxy
GALEX0332-2823	53.240459	-28.388306	20.83	...	0.2191	0.2137	79± 0	Galaxy
GALEX0333-2821	53.258461	-28.357668	21.08	...	0.2507	0.2472	95± 1	Galaxy
GALEX0334-2815	53.541245	-28.255445	21.57	...	0.3450	0.3356	2± 0	Galaxy
GALEX0959+0149	149.873642	1.833014	21.42	20.10	0.2076	0.2052	30± 0	Galaxy
GALEX0959+0151	149.917755	1.855986	21.10	20.42	0.2536	0.2507	1019± 2	Galaxy
GALEX1000+0201	150.148819	2.020419	20.71	20.11	0.2709	0.2653	247± 1	Galaxy
GALEX1001+0233	150.413162	2.563489	21.00	20.90	0.3889	0.3823	435± 2	Galaxy
GALEX1047+5827	161.772766	58.461445	20.33	19.27	0.2479	0.2439	... 1	Seyfert 1.8
GALEX1240+6233	190.041367	62.561611	20.28	19.33	0.2104	0.2068	164± 1	Galaxy
GALEX1417+5232	214.428314	52.538250	21.46	22.08	0.1989	0.1923	59± 1	Galaxy
GALEX1417+5228	214.431000	52.468281	21.90	19.30	0.2104	0.2067	1390± 2	Galaxy
GALEX1417+5242	214.474167	52.700405	21.71	20.95	0.2047	0.2035	33± 1	Galaxy
GALEX1418+5307	214.728745	53.129887	22.09	21.99	0.2047	0.2035	357± 3	Galaxy
GALEX1418+5259	214.732956	52.992176	21.38	21.15	0.2882	0.2871	194± 1	Galaxy
GALEX1419+5223	214.811447	52.390835	21.77	20.69	0.2565	0.2542	85± 6	Galaxy
GALEX1419+5230	214.907440	52.506832	21.18	20.03	0.2853	0.2815	13± 1	Galaxy
GALEX1420+5247	215.133041	52.799362	20.73	20.37	0.2565	0.2515	93± 17	Galaxy
GALEX1420+5250	215.186127	52.835140	21.36	20.47	0.2536	0.2519	80± 3	Galaxy
GALEX1421+5239	215.352615	52.655499	21.82	20.06	0.2565	0.2594	42± 1	Galaxy
GALEX1422+5252	215.533951	52.873802	21.30	19.60	0.3055	0.3017	12± 0	Seyfert 1.5
GALEX1436+3456	219.092209	34.942055	21.16	20.65	0.2716	0.2684	88± 4	Galaxy
GALEX1436+3459	219.100418	34.993500	21.31	...	0.2148	0.2115	73± 1	Galaxy
GALEX1436+3525	219.193085	35.417583	21.22	19.75	0.3697	0.3688	7± 0	Seyfert 1.8
GALEX1437+3445	219.333954	34.757584	21.18	20.84	0.3285	0.3237	182± 1	Galaxy
GALEX1437+3522	219.388123	35.370167	20.64	20.78	0.2475	0.2428	16± 1	Galaxy
GALEX1712+5956	258.149261	59.946831	20.37	20.42	0.2248	0.2206	522± 4	Galaxy
GALEX1712+6001	258.197540	60.022892	20.80	20.97	0.7863	0.7780	...	Galaxy
GALEX1714+5950	258.591827	59.833359	21.09	20.25	0.2306	0.2298	104± 1	Galaxy
GALEX1714+5956	258.684052	59.947475	21.13	20.49	0.2220	0.2156	33± 1	Galaxy
GALEX1717+5944	259.465027	59.748306	21.08	20.05	0.1960	0.1979	135± 1	Galaxy

TABLE 12
DEIMOS OBSERVATIONS: *GALEX* AGNS

NAME (1)	R.A. (J2000.0) (2)	Decl. (J2000.0) (3)	NUV (4)	z_{galex} (5)	$z_{optical}$ (6)	Optical Type (7)
GALEX0332-2810	53.184074	-28.174442	21.04	1.1519	1.1480	AGN
GALEX1000+0152	150.139114	1.876969	21.08	0.8323	0.8309	AGN
GALEX1000+0202	150.145584	2.043078	21.25	1.1778	1.1764	AGN
GALEX1417+5230	214.399780	52.508324	20.82	0.9878	0.9851	AGN
GALEX1418+5237	214.519180	52.629162	21.32	1.0900	1.0885	AGN
GALEX1418+5310	214.657745	53.171547	21.01	0.8957	0.8922	AGN
GALEX1420+5240	215.042847	52.674995	21.17	0.5526	0.5483	Galaxy
GALEX1436+3453	219.131790	34.890221	20.33	1.0677	1.0652	AGN
GALEX1436+3450	219.152878	34.842777	20.66	0.2830	0.2810	AGN
GALEX1437+3522	219.311295	35.381779	19.33	0.6299	0.6236	AGN
GALEX1437+3448	219.345505	34.801777	20.98	0.9497	0.9476	AGN
GALEX1437+3445	219.370041	34.763195	19.22	0.6284	0.6224	AGN
GALEX1712+6007	258.131622	60.122139	20.98	1.2037	1.2020	AGN
GALEX1712+6005	258.145172	60.091831	19.55	0.9993	0.9973	AGN
GALEX1715+5956	258.751862	59.936611	19.71	0.6950	0.7000	AGN

TABLE 13
DEIMOS OBSERVATIONS: *GALEX* SPECTRA WITHOUT STRONG $\text{Ly}\alpha$ EMISSION

NAME (1)	R.A. (J2000.0) (2)	Decl. (J2000.0) (3)	NUV (4)	g (5)	$z_{optical}$ (6)	EW($\text{H}\alpha$) (7)	Optical Type (8)
GALEX0331-2810	52.932369	-28.173195	21.21	...	0.2169	21± 1	Galaxy
GALEX0331-2811	52.948586	-28.193085	21.46	...	0.3814	81± 8	Galaxy
GALEX0331-2810	52.949966	-28.168806	21.19	...	0.3798	56± 1	Galaxy
GALEX0331-2811	52.952209	-28.184692	20.67	...	0.2157	103± 1	Galaxy
GALEX0331-2812	52.953033	-28.208719	20.79	...	0.2136	31± 1	Galaxy
GALEX0331-2813	52.958126	-28.221001	21.27	...	0.2730	93± 3	Galaxy
GALEX0331-2813	52.968121	-28.231945	21.04	...	0.2793	37± 1	Galaxy
GALEX0331-2814	52.978458	-28.235889	21.26	...	0.3160	81± 1	Galaxy
GALEX0331-2812	52.980709	-28.208830	21.46	...	0.2775	9± 0	Galaxy
GALEX0332-2805	53.022079	-28.096029	21.45	...	0.3866	76± 4	Galaxy
GALEX0332-2808	53.026207	-28.134026	21.55	...	0.2775	54± 1	Galaxy
GALEX0332-2810	53.155334	-28.177498	21.01	...	0.2035	20± 0	Galaxy
GALEX0332-2810	53.168877	-28.182112	21.27	...	0.1955	72± 1	Galaxy
GALEX0333-2813	53.421875	-28.223223	21.19	...	0.3593	60± 1	Galaxy
GALEX0333-2819	53.426876	-28.322580	21.24	...	0.2332	56± 1	Galaxy
GALEX0333-2821	53.448418	-28.364719	20.58	...	0.2480	15± 1	Galaxy
GALEX0331-2810	52.932369	-28.173195	21.21	-99.00	0.2169	21± 1	Galaxy
GALEX0331-2811	52.948586	-28.193085	21.46	-99.00	0.3814	81± 8	Galaxy
GALEX0331-2810	52.949966	-28.168806	21.19	-99.00	0.3798	56± 0	Galaxy
GALEX0331-2811	52.952209	-28.184692	20.67	-99.00	0.2157	103± 1	Galaxy
GALEX0331-2812	52.953033	-28.208719	20.79	-99.00	0.2136	31± 1	Galaxy
GALEX0331-2813	52.958126	-28.221001	21.27	-99.00	0.2730	93± 3	Galaxy
GALEX0331-2813	52.968121	-28.231945	21.04	-99.00	0.2793	37± 1	Galaxy
GALEX0331-2814	52.978458	-28.235889	21.26	-99.00	0.3160	81± 1	Galaxy
GALEX0331-2812	52.980709	-28.208830	21.46	-99.00	0.2775	9± 0	Galaxy
GALEX0332-2805	53.022079	-28.096029	21.45	-99.00	0.3866	76± 4	Galaxy
GALEX0332-2808	53.026207	-28.134026	21.55	-99.00	0.2775	54± 1	Galaxy
GALEX0332-2810	53.155334	-28.177498	21.01	-99.00	0.2035	20± 0	Galaxy
GALEX0332-2810	53.168877	-28.182112	21.27	-99.00	0.1955	72± 1	Galaxy
GALEX0333-2813	53.421875	-28.223223	21.19	-99.00	0.3593	60± 1	Galaxy
GALEX0333-2819	53.426876	-28.322580	21.24	-99.00	0.2332	56± 1	Galaxy
GALEX0333-2821	53.448418	-28.364719	20.58	-99.00	0.2480	15± 1	Galaxy
GALEX1417+5240	214.465881	52.679333	21.55	22.99	0.3666	84± 2	Galaxy
GALEX1417+5237	214.490631	52.632240	20.94	19.65	0.2849	19± 1	Galaxy
GALEX1417+5228	214.493362	52.482773	21.44	20.44	0.3710	50± 2	Galaxy
GALEX1418+5309	214.686539	53.166164	21.52	21.35	0.2578	57± 1	Galaxy
GALEX1418+5303	214.744461	53.060051	21.78	20.55	0.2045	32± 1	Galaxy
GALEX1419+5229	214.889496	52.489220	21.78	20.46	0.2464	33± 3	Galaxy
GALEX1420+5243	215.099930	52.726917	21.46	19.76	0.2396	35± 1	Galaxy
GALEX1420+5246	215.162506	52.775528	21.81	20.05	0.2397	20± 1	Galaxy
GALEX1420+5251	215.203964	52.864445	21.73	20.75	0.2897	38± 1	Galaxy
GALEX1422+5246	215.514145	52.768780	21.78	21.28	0.4179	3± 1	Galaxy
GALEX1422+5251	215.540207	52.859055	21.59	20.95	0.3808	102± 2	Galaxy
GALEX1436+3457	219.150085	34.958279	21.22	19.73	0.2505	17± 2	Galaxy
GALEX1712+5955	258.151184	59.930222	20.76	20.93	0.2720	138± 1	Galaxy
GALEX1718+5945	259.503265	59.755474	20.81	18.90	0.2290	5± 0	Galaxy
GALEX1718+5944	259.533936	59.748917	20.89	19.45	0.3899	52± 3	Galaxy

TABLE 14
GALEX X-RAY PROPERTIES

NAME (1)	R.A. (J2000.0) (2)	Decl. (J2000.0) (3)	$f_{2-8 \text{ keV}}$ (4)	$f_{0.5-2 \text{ keV}}$ (5)	$\log(L_{2-8 \text{ keV}})$ (6)	z_{galax} (7)	class (8)
GALEX0332-2740	53.1104	-27.6766	1.54e-14	2.20e-14	43.91	1.0396	AGN
GALEX0332-2741	53.1125	-27.6847	6.55e-14	5.97e-14	44.19	0.7373	AGN
GALEX0331-2756	52.8365	-27.9469	1.70e-14	1.03e-14	43.53	0.6850	AGN
GALEX0332-2739	53.1588	-27.6625	2.55e-14	2.34e-14	43.91	0.8381	AGN
GALEX0333-2756	53.3705	-27.9447	4.98e-14	3.91e-14	44.20	0.8381	AGN
GALEX1048+5910	162.049	59.1797	2.23e-14	2.58e-14	43.68	0.7110	AGN
GALEX1049+5853	162.372	58.8940	8.69e-14	9.08e-14	43.24	0.2506	AGN
GALEX1237+6217	189.279	62.2840	1.89e-14	1.57e-14	43.98	1.019	AGN
GALEX1237+6211	189.498	62.1840	9.11e-14	5.64e-14	44.55	0.9129	AGN
GALEX0332-2753	53.2362	-27.8879	-4.67e-16	2.05e-16	< 41.0	0.3745	Galaxy
GALEX0332-2801	53.0497	-28.0250	...	...	< 40.5	0.2191	Galaxy

TABLE 15
LOW-REDSHIFT GALEX LAE SAMPLE

NAME (1)	R.A. (J2000.0) (2)	Decl. (J2000.0) (3)	NUV (4)	FUV (5)	z_{galax} (6)	$\log L(\text{Ly}\alpha)$ (7)	EW(Ly α) (8)	z_{optical} (9)
GALEX1718+5949	259.748260	59.825752	20.91	23.31	0.3717	42.24 $\pm$ 0.08	115 $\pm$ 49	...
GALEX1417+5228	214.430923	52.468307	21.90	21.79	0.2104	42.01 $\pm$ 0.01	101 $\pm$ 9	0.2067 SF
GALEX0331-2814	52.976460	-28.238554	21.49	21.84	0.2824	42.15 $\pm$ 0.02	84 $\pm$ 9	0.2802 SF
GALEX1436+3525	219.193008	35.417641	21.22	22.20	0.3697	42.32 $\pm$ 0.05	81 $\pm$ 16	0.3689 AGN
GALEX0334-2815	53.541370	-28.255472	21.57	22.77	0.3450	41.93 $\pm$ 0.08	75 $\pm$ 27	0.3348 AGN
GALEX1422+5252	215.533752	52.873722	21.30	22.52	0.3055	42.01 $\pm$ 0.02	71 $\pm$ 8	0.3017 AGN
GALEX0332-2801	53.049873	-28.025000	21.51	21.71	0.2191	41.76 $\pm$ 0.02	67 $\pm$ 9	0.2155 SF
GALEX1240+6233	190.041428	62.561668	20.28	20.69	0.2104	42.01 $\pm$ 0.04	61 $\pm$ 10	0.2068 BPT AGN
GALEX1437+3500	219.406250	35.011749	21.89	23.15	0.3853	42.07 $\pm$ 0.10	59 $\pm$ 18	...
GALEX1418+5307	214.728928	53.130028	22.09	21.95	0.2047	41.73 $\pm$ 0.02	53 $\pm$ 6	0.2035 SF
GALEX1421+5249	215.463165	52.830891	21.25	21.87	0.2018	41.50 $\pm$ 0.06	49 $\pm$ 9	...
GALEX1438+3457	219.746078	34.960636	21.29	21.93	0.3740	42.27 $\pm$ 0.07	48 $\pm$ 10	...
GALEX1419+5315	214.794739	53.265835	20.98	21.90	0.2651	41.85 $\pm$ 0.03	48 $\pm$ 7	...
GALEX1001+0220	150.294846	2.346972	21.47	21.51	0.2508	42.05 $\pm$ 0.01	45 $\pm$ 3	(0.2481)pr
GALEX1000+0157	150.115784	1.951139	20.72	20.69	0.2680	42.42 $\pm$ 0.01	43 $\pm$ 2	(0.2645)pr
GALEX1435+3449	218.764587	34.821220	21.32	22.75	0.3782	42.03 $\pm$ 0.12	42 $\pm$ 14	...
GALEX1714+5949	258.591766	59.833332	21.09	21.28	0.2306	41.94 $\pm$ 0.03	38 $\pm$ 5	0.2298 SF
GALEX0332-2811	53.174255	-28.190195	20.27	20.17	0.2076	42.15 $\pm$ 0.01	38 $\pm$ 2	0.2044 SF
GALEX0331-2751	52.759998	-27.858416	21.50	21.97	0.3400	42.02 $\pm$ 0.05	35 $\pm$ 7	...
GALEX1436+3459	219.100540	34.993530	21.31	21.57	0.2148	41.73 $\pm$ 0.03	34 $\pm$ 4	0.2115 SF
GALEX1418+5259	214.733200	52.992195	21.38	21.88	0.2882	41.80 $\pm$ 0.04	34 $\pm$ 5	0.2871 SF
GALEX0038-4352	9.520541	-43.874584	20.63	21.08	0.2220	41.89 $\pm$ 0.03	33 $\pm$ 4	0.2191 SF
GALEX0332-2823	53.240330	-28.388361	20.83	21.30	0.2191	41.70 $\pm$ 0.03	33 $\pm$ 4	0.2137 SF
GALEX1437+3445	219.334000	34.757637	21.18	21.53	0.3285	42.23 $\pm$ 0.03	32 $\pm$ 3	0.3237 SF
GALEX1436+3440	219.086914	34.672001	21.09	22.04	0.3740	42.08 $\pm$ 0.12	32 $\pm$ 9	...
GALEX1001+0233	150.413330	2.563611	21.00	21.43	0.3889	42.42 $\pm$ 0.06	32 $\pm$ 5	0.3823 SF
GALEX1417+5246	214.438126	52.771694	21.83	22.09	0.2479	41.57 $\pm$ 0.05	32 $\pm$ 6	...
GALEX0332-2811	53.061497	-28.186584	21.18	21.71	0.2651	41.68 $\pm$ 0.05	30 $\pm$ 5	0.2789 SF ?
GALEX1437+3441	219.356308	34.685585	21.00	21.64	0.2944	41.96 $\pm$ 0.05	28 $\pm$ 5	...
GALEX1417+5305	214.291290	53.086723	21.29	21.85	0.2680	41.65 $\pm$ 0.04	26 $\pm$ 4	...
GALEX1418+5218	214.593506	52.306747	21.70	22.35	0.2392	41.44 $\pm$ 0.05	26 $\pm$ 5	...
GALEX1434+3532	218.718338	35.547527	21.44	21.81	0.1977	41.45 $\pm$ 0.07	25 $\pm$ 5	...
GALEX1436+3456	219.092041	34.942139	21.16	21.47	0.2716	41.91 $\pm$ 0.03	25 $\pm$ 3	0.2684 SF
GALEX0959+0149	149.873871	1.832972	21.42	21.76	0.2076	41.27 $\pm$ 0.10	25 $\pm$ 7	0.2052 SF
GALEX1423+5246	215.775833	52.779697	21.33	22.01	0.3458	41.86 $\pm$ 0.06	23 $\pm$ 4	...
GALEX0331-2811	52.962204	-28.189028	20.68	21.02	0.2162	41.64 $\pm$ 0.03	23 $\pm$ 3	0.2132 SF
GALEX0959+0151	149.917953	1.856028	21.10	21.47	0.2536	41.77 $\pm$ 0.05	22 $\pm$ 3	0.2507 SF
GALEX1420+5306	215.170456	53.114002	21.05	21.49	0.1989	41.32 $\pm$ 0.08	22 $\pm$ 5	...
GALEX1421+5239	215.352737	52.655472	21.82	22.27	0.2565	41.48 $\pm$ 0.05	22 $\pm$ 4	(0.2578)f AGN
GALEX0330-2816	52.737583	-28.279501	21.11	21.55	0.2853	41.80 $\pm$ 0.05	22 $\pm$ 4	...
GALEX1714+5956	258.683746	59.947247	21.13	21.62	0.2220	41.53 $\pm$ 0.10	21 $\pm$ 6	0.2156 SF
GALEX1420+5247	215.132080	52.799389	20.73	21.13	0.2565	41.83 $\pm$ 0.02	21 $\pm$ 2	0.2515 SF
GALEX1725+5920	261.344940	59.345528	20.01	20.35	0.3861	42.43 $\pm$ 0.14	20 $\pm$ 7	...
GALEX1717+5944	259.465027	59.748554	21.08	21.66	0.1960	41.40 $\pm$ 0.13	20 $\pm$ 6	0.1979 SF
GALEX1716+5932	259.178070	59.548889	21.04	21.79	0.3600	41.99 $\pm$ 0.14	20 $\pm$ 7	...
GALEX1001+0145	150.396637	1.756500	21.50	21.74	0.2191	41.40 $\pm$ 0.08	20 $\pm$ 4	...
GALEX1423+5244	215.842850	52.742474	21.74	22.36	0.2795	41.46 $\pm$ 0.09	20 $\pm$ 5	...
GALEX1421+5224	215.477295	52.406723	20.39	21.22	0.3515	42.12 $\pm$ 0.04	20 $\pm$ 2	...
GALEX0333-2821	53.258461	-28.357695	21.08	21.26	0.2507	41.64 $\pm$ 0.05	20 $\pm$ 3	0.2472 SF
GALEX1000+0201	150.149094	2.020583	20.71	21.03	0.2709	41.99 $\pm$ 0.03	19 $\pm$ 2	0.2653 SF
GALEX1418+5245	214.521545	52.752110	21.30	21.63	0.2450	41.60 $\pm$ 0.04	19 $\pm$ 3	...
GALEX1419+5223	214.811249	52.390808	21.77	22.16	0.2565	41.35 $\pm$ 0.08	19 $\pm$ 4	0.2542 SF
GALEX1419+5221	214.970749	52.350250	21.29	21.83	0.2680	41.57 $\pm$ 0.06	19 $\pm$ 4	...
GALEX1418+5217	214.701202	52.299168	21.66	22.03	0.2421	41.42 $\pm$ 0.07	19 $\pm$ 4	...
GALEX0331-2809	52.999332	-28.164444	21.18	21.43	0.2392	41.51 $\pm$ 0.05	19 $\pm$ 3	0.2364 SF
GALEX1717+5929	259.379578	59.487473	20.34	20.94	0.2997	41.99 $\pm$ 0.07	18 $\pm$ 4	...
GALEX1418+5249	214.732056	52.824558	21.16	21.13	0.2680	41.48 $\pm$ 0.07	18 $\pm$ 4	(0.2630)f AGN
GALEX1420+5243	215.180359	52.718750	21.29	21.59	0.2508	41.63 $\pm$ 0.04	18 $\pm$ 2	...
GALEX1436+3527	219.026245	35.458553	21.38	21.57	0.2517	41.79 $\pm$ 0.04	17 $\pm$ 2	...
GALEX1422+5310	215.500092	53.168472	21.65	22.28	0.2392	41.17 $\pm$ 0.12	17 $\pm$ 5	...
GALEX1420+5250	215.186218	52.835056	21.36	21.64	0.2536	41.58 $\pm$ 0.04	17 $\pm$ 2	0.2519 SF
GALEX1419+5230	214.907211	52.507057	21.28	21.75	0.2853	41.55 $\pm$ 0.07	16 $\pm$ 3	0.2815 SF

TABLE 16
MODERATE-REDSHIFT *GALEX* LAE SAMPLE

NAME (1)	R.A. (J2000.0) (2)	Decl. (J2000.0) (3)	NUV (4)	FUV (5)	z_{galax} (6)	$\log L(\text{Ly}\alpha)$ (7)	EW(Ly α) (8)	$z_{optical}$ (9)
GALEX1421+5221	215.421539	52.359886	21.64	25.20	0.9475	43.21 $\pm$ 0.02	52 $\pm$ 4	...
GALEX1712+6001	258.197723	60.022778	20.80	23.63	0.7863	43.19 $\pm$ 0.03	37 $\pm$ 4	0.7780 SF
GALEX1437+3541	219.281967	35.692280	20.93	22.79	0.6640	42.91 $\pm$ 0.05	30 $\pm$ 5	...
GALEX1420+5313	215.237045	53.218750	21.26	23.87	0.7488	42.84 $\pm$ 0.04	27 $\pm$ 3	...